

Disclosure: E.B. Johannsen: None. A. Berglund: None. A. Skakkebaek: None. S. Chang: None. J. Rohayem: None. S. Laurentino: None. A. Hørlyck: None. S.O. Drue: None. E.N. Bak: None. J. Fedder: None. F. Tuttmann: None. J. Gromoll: None. J. Just: None. C.H. Gravholt: None.

Background: 46,XX testicular disorder of sex development is a rare congenital condition affecting approximately three per 100,000 newborn males. It is characterized by a combination of the typical female sex chromosome constitution, 46,XX, and a variable male phenotype. In the majority of XX males, a Y chromosome segment containing the sex-determining region gene (*SRY*) has been translocated to the paternal X chromosome. However, the precise genomic content of the translocated segment, breakpoints, genome-wide effects, and the effect on other biological pathways essential for the phenotype remain unknown. **Methods:** We performed long-read DNA sequencing (Oxford Nanopore), RNA sequencing (paired-end, 100 bp) and DNA methylation (Infinium EPIC) analysis on blood samples from males with 46,XX testicular disorder of sex development (XX males; n = 11), male controls (46,XY; n = 12 for long-read DNA sequencing, n = 26 for RNA sequencing and DNA methylation) and female controls (46,XX; n = 12 for long-read DNA sequencing, n = 32 for RNA sequencing and DNA methylation), in addition to RNA sequencing and DNA methylation analysis on blood samples from males with Klinefelter syndrome (47,XXY, n = 22). We also performed clinical measures on all XX males and a subset of male controls (n = 10). **Results:** We identified variation in the translocated Y-chromosome segment of our XX male cohort, allowing for subcategorization into 1) XX males without Y chromosome material (n = 1), 2) those with short Yp arms (breakpoint at 2.7-2.8 Mbp, n = 2), 3) those with medium Yp arms (breakpoint at 7.3 Mbp, n = 1) and 4) those with long Yp arms, including *AMELY*, *TBLY1* deletions, and in some cases *PRKY* deletions (n = 7). We also identified variable expression of the X-Y homologues *PRKY* and *PRKX*, appeared to have balanced expression level resulting in overall equilibrium. The Y-chromosomal transcriptome and methylome were affected, reflecting the Y-chromosome segment length. Genomic changes induced by the translocated Yp segment were evident in the transcriptome and methylome of autosomal and X-chromosomal regions, indicating global effects. Furthermore, the transcriptional changes in XX males correlated with phenotypic traits of the XX males, including lower height, decreased lean mass and smaller testicular size. **Conclusion:** Integration of long-read DNA sequencing, DNA methylation, and RNA sequencing analysis allowed for the identification of a heterogeneous translocation process in XX males that exerted widespread effects on methylation and transcription.

Presentation: 6/2/2024

Abstract citation ID: bvae163.1621

Reproductive Endocrinology 8320

Integration of long-read sequencing, DNA methylation and gene expression reveals heterogeneity in Y-chromosome segment lengths in phenotypic males with 46,XX testicular disorder of sex development

Emma Bruun Johannsen, MSc¹, Agnethe Berglund, MD, PhD¹, Anne Skakkebaek, MD, PhD¹, Simon Chang¹, Julia Rohayem, MD, PhD², Sandra Laurentino, PhD², Arne Hørlyck, MD¹, Simon Opstrup Drue, PhD¹, Ebbe Norskov Bak, PhD¹, Jens Fedder, MD, PhD³, Frank Tuttmann⁴, Joerg Gromoll, PhD², Jesper Just, PhD⁵, and Claus Hojbjerg Gravholt, MD, PhD¹

¹Aarhus University Hospital, Aarhus, Denmark; ²Münster University Hospital, Münster, Germany; ³Odense University Hospital, Odense, Denmark; ⁴University of Münster, Münster, Germany; ⁵Aarhus University Hospital, Aarhus N, Denmark