

to bias and had some limitations. Historic vaccine uptake is low, which may diminish the impact of maternal vaccination.

This study demonstrates meaningful benefit of COVID-19 vaccination during pregnancy, which has moderately waned over time, but still supports the need to reconsider the decision to remove vaccine eligibility during pregnancy. Continued studies on the COVID-19 risk and vaccine effectiveness in this population are important.

Nathan C. Lo, MD, PhD
Yvonne Maldonado, MD
Helen Y. Chu, MD, MPH
Peter J. Hotez, MD, PhD
Mathew V. Kiang, ScD

Author Affiliations: Division of Infectious Diseases and Geographic Medicine, Department of Medicine, Stanford University, Stanford, California (Lo); Departments of Pediatrics and Epidemiology and Population Health, Stanford University, Stanford, California (Maldonado); Division of Allergy and Infectious Diseases, Department of Medicine, University of Washington, Seattle (Chu); Texas Children's Hospital Center for Vaccine Development, Departments of Pediatrics and Molecular Virology and Microbiology, National School of Tropical Medicine, Baylor College of Medicine, Houston, Texas (Hotez); Department of Epidemiology and Population Health, Stanford University, Stanford, California (Kiang).

Accepted for Publication: July 25, 2025.

Published Online: September 29, 2025. doi:10.1001/jamapediatrics.2025.3561

Correction: This article was corrected on October 27, 2025, to correct the panel label of Figure 2B.

Corresponding Author: Nathan C. Lo, MD, PhD, Division of Infectious Diseases and Geographic Medicine, Stanford University, Stanford, CA 94304 (nathan.lo@stanford.edu).

Author Contributions: Drs Lo and Kiang had full access to all of the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.

Concept and design: Lo, Kiang.

Acquisition, analysis, or interpretation of data: All authors.

Drafting of the manuscript: Lo, Kiang.

Critical review of the manuscript for important intellectual content: All authors.

Statistical analysis: Lo, Kiang.

Supervision: Lo, Kiang.

Conflict of Interest Disclosures: Dr Lo reported personal fees from the World Health Organization related to guidelines on neglected tropical diseases outside the submitted work. Dr Maldonado reported grants from Pfizer outside the submitted work. Dr Chu reported personal fees from AbbVie, Merck, Roche, and Vir Biotechnology outside the submitted work. Dr Hotez reported being a coinventor on non-revenue-generating patents for neglected tropical diseases, which are owned by Baylor College of Medicine, and a coinventor of a COVID-19 recombinant protein vaccine technology, owned by Baylor College of Medicine, which was licensed by Baylor Ventures nonexclusively and without patent restrictions to several companies committed to advancing vaccine technology for low- and middle-income countries, and receiving royalty income from American Society for Microbiology-Wiley and Johns Hopkins University Press. Dr Kiang reported grants from the US National Institutes of Health outside the submitted work.

Data Sharing Statement: See Supplement 2.

1. Simeone RM, Zambrano LD, Halasa NB, et al; Overcoming COVID-19 Investigators. Effectiveness of maternal mRNA COVID-19 vaccination during pregnancy against COVID-19-associated hospitalizations in infants aged <6 months during SARS-CoV-2 Omicron predominance - 20 states, March 9, 2022-May 31, 2023. *MMWR Morb Mortal Wkly Rep.* 2023;72(39):1057-1064. doi:10.15585/mmwr.mm7239a3

2. US Centers for Disease Control and Prevention. COVID-NET. Accessed June 11, 2025. <https://www.cdc.gov/covid/php/covid-net/index.html>

3. McClymont E, Albert AY, Alton GD, et al; CANCOVID-Preg Team. Association of SARS-CoV-2 infection during pregnancy with maternal and perinatal outcomes. *JAMA.* 2022;327(20):1983-1991. doi:10.1001/jama.2022.5906

4. Link-Gelles R, Chickery S, Webber A, et al; CDC COVID-19 Vaccine Effectiveness Collaborators. Interim estimates of 2024-2025 COVID-19 vaccine effectiveness among adults aged ≥18 years - VISION and IVY Networks, September 2024-January 2025. *MMWR Morb Mortal Wkly Rep.* 2025;74(6):73-82. doi:10.15585/mmwr.mm7406a1

5. Guedalia J, Lipschuetz M, Calderon-Margalit R, et al. Effectiveness of a third BNT162b2 mRNA COVID-19 vaccination during pregnancy: a national observational study in Israel. *Nat Commun.* 2022;13(1):6961. doi:10.1038/s41467-022-34605-x

6. Magnus MC, Örtqvist AK, Dahlqvist E, et al. Association of SARS-CoV-2 vaccination during pregnancy with pregnancy outcomes. *JAMA.* 2022;327(15):1469-1477. doi:10.1001/jama.2022.3271

Clinical Long-Read Sequencing Test for Genetic Disease Diagnosis

Recent work in genome sequencing (GS) shows a 4% to 8% increase in diagnostic yield over exome sequencing (ES) alone, albeit the overall diagnostic gain (<1%) is marginal if GS is compared to combinations of ES with other standard tests.¹



Supplemental content

Recently, we reported undiagnosed cases following GS could be recovered by long-read sequencing (LRS), due in part to the simultaneous detection of clinically relevant repeat expansions,²⁻⁴ methylation abnormalities,⁵ and improved structural variant detection.⁶ However, no study has assessed the implementation of LRS as a first-line clinical test. Herein, we report the comparative diagnostic yield and turnaround time and evaluate the ability to consolidate standard-of-care (SOC) approaches using a single clinical LRS test.

Methods | Clinical high-fidelity (HiFi) LRS was performed on 235 patients aged 0 to 18 years compared to an age- and phenotype-matched control cohort of 513 patients. Patients selected for the control group underwent SOC inpatient genetic testing, including expedited ES/GS, karyotype, fluorescence in situ hybridization (FISH), chromosomal microarray (CMA), and targeted panels. The eMethods in Supplement 1 provide further details. The Standards for Reporting of Diagnostic Accuracy (STARD) reporting guidelines were followed.

Results | LRS case results were compared to controls (Figure 1), with the true diagnostic rate determined by counting diagnoses explaining the primary signs and symptoms. Variants of uncertain significance, carrier status, incidental findings, or pathogenic changes associated with late-onset or variable phenotypes not directly linked to presenting complaint were excluded.

The LRS cases yielded a significantly higher ($P = .004$, 2-2 χ^2 test) diagnostic rate compared to the SOC group (Figure 2). The mean length of time from test order to diagnostic report (27 days) was significantly ($P = .048$, t test) shorter among LRS cases (Figure 2) compared to controls (62 days). Similarly, if a negative result was returned, the average time to report was significantly ($P < .001$, t test) shorter among LRS cases (29 days) compared to controls (91 days).

Factors negatively impacting the time to diagnosis specific to LRS were low sequencing yields requiring reloading and technical limitations of tertiary analysis software. Among the SOC group, the primary cause for delayed diagnosis was mul-

Figure 1. Diagram Reporting Flow of Probands and Number of Test Orders in Long-Read Sequencing (LRS) and Standard-of-Care (SOC) Cohorts





Figure 2. Long-Read Sequencing (LRS) Compared to Standard Genetic Testing Among Matched Pediatric Cases



P values calculated using Fisher exact test. CMA indicates chromosomal microarray analysis; ES, exome sequencing; GS, genome sequencing.

multiple test orders required to reach a diagnosis, with an average of 6.1 tests required in the SOC group compared to 2.7 tests in LRS cases. When limited to diagnostic cases, SOC controls had an average of 1.6 test orders for each diagnostic finding ($n = 138$), compared to 1.01 among LRS cases ($n = 87$, with 1 diagnosis requiring follow-up cytogenetic testing). If the first test was positive in controls, the mean time to diagnosis was comparable to LRS. However, 57 SOC cases required multiple tests to reach a diagnosis in which the average turnaround time was more than 4 months.

A large fraction of variation in LRS originated from complex structural variants (SVs) and copy number variants, repeat expansions, and methylation variation. Specifically, 16 of 87 reported diagnostic variants in the LRS group (18.3%) benefited from the integrated capability of LRS, which included aberrant methylation, rare expansion disorders, phasing of single-nucleotide variation in a singleton, and detection or refinement of SVs. Note that the availability of standard clinical testing is limited, such as *GNAS* methylation and newly described repeat expansions (eg, in *DIP2B*), which were diagnostic findings captured by LRS. As expected, 71 of 87 diagnostic genotypes (81.6%) without previous standard genetic testing would have been detected by ES, GS, or CMA.

Discussion | Clinical LRS offers a single, comprehensive genomic assay for diagnosing genetic disease. The advantage of LRS was explained by its expanded variant detection. While

the framework used to interpret LRS variation relies on changes also detected by combinations of standard tests, the difference is that in clinical practice, all testing modalities are not typically deployed, and some tests, such as repeat expansions, may not be included in the differential if a presentation is atypical. Due to the sequential nature of SOC testing, there are significant delays in reaching diagnoses compared to LRS. These benefits of clinical LRS are likely only the beginnings of what may be uncovered through expanded testing and further understanding of noncoding regions.^{5,6}

Isabelle Thiffault, PhD
Emily Farrow, PhD
Cassandra Barrett, PhD
Meadow Scott, BSc
Amy Ross, BSc
John C. Means, PhD
Warren A. Cheung, PhD
Adam F. Johnson, PhD
Boryana Koseva, PhD
Rebecca McLennan, PhD
Elin Grundberg, PhD
Chengpeng Bi, PhD
Carl Schwendinger-Schreck, PhD
Byunggil Yoo, MSc
Jeffrey J. Johnston, BSc
Florescia Del Viso, PhD

Vitoria Paolillo, PhD
 John Herriges, PhD
 Lei Zhang, PhD
 Margaret Gibson, BSc
 Ana S. A. Cohen, PhD
 Joe Alaimo, PhD
 Carol J. Saunders, PhD
 Tomi Pastinen, MD, PhD

Author Affiliations: Department of Pathology and Laboratory Medicine, Children's Mercy Kansas City, Kansas City, Missouri (Thiffault, Farrow, Del Viso, Paolillo, Herriges, Zhang, Gibson, Cohen, Alaimo, Saunders); Division of Clinical Genetics, Department of Pediatrics, Children's Mercy Kansas City, Kansas City, Missouri (Barrett); Genomic Medicine Center, Department of Pediatrics, Children's Mercy Kansas City, Kansas City, Missouri (Scott, Ross, Means, Cheung, Johnson, Koseva, McLennan, Grundberg, Bi, Schwendinger-Schreck, Yoo, Johnston, Pastinen).

Accepted for Publication: July 23, 2025.

Published Online: September 22, 2025. doi:10.1001/jamapediatrics.2025.3320

Open Access: This is an open access article distributed under the terms of the CC-BY License. © 2025 Thiffault I et al. JAMA Pediatrics.

Corresponding Author: Tomi Pastinen, MD, PhD, Children's Mercy Kansas City, 2401 Gillham Rd, Kansas City, MO 64108 (tpastinen@cmh.edu).

Author Contributions: Drs Thiffault and Pastinen had full access to all of the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis. Drs Thiffault and Farrow contributed equally. *Concept and design:* Thiffault, Farrow, McLennan, Saunders, Pastinen.

Acquisition, analysis, or interpretation of data: Thiffault, Farrow, Barrett, Scott, Ross, Means, Cheung, Johnson, Koseva, Grundberg, Bi, Schwendinger-Schreck, Yoo, Johnston, Del Viso, Paolillo, Herriges, Gibson, Cohen, Alaimo, Saunders, Pastinen.

Drafting of the manuscript: Thiffault, Farrow, McLennan, Saunders, Pastinen.

Critical review of the manuscript for important intellectual content: All authors.

Statistical analysis: Johnson, Koseva, Grundberg, Schwendinger-Schreck, Pastinen.

Obtained funding: Pastinen.

Administrative, technical, or material support: Barrett, Scott, Ross, Cheung, McLennan, Grundberg, Bi, Yoo, Gibson.

Supervision: Thiffault, Grundberg, Gibson, Saunders, Pastinen.

Conflict of Interest Disclosures: Dr Pastinen and his team reported nonfinancial support from Pacific Biosciences, including reagent support and speaker travel fees, during the conduct of the study and serving on the scientific advisory board for Hitachi High-Tech; and operating grant (R01) support from the US National Institutes of Health outside the submitted work. No other disclosures were reported.

Funding/Support: Genomic Answers for Kids provided salary support for research infrastructure and staff. Pacific Biosciences provided in-kind contribution for the high-fidelity genome sequencing reagents.

Role of the Funder/Sponsor: The funders had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

Data Sharing Statement: See Supplement 2.

Additional Contributions: The Genomic Answers for Kids program is made possible by the generous gifts to Children's Mercy Research Institute at Children's Mercy Kansas City.

Additional Information: Institutional review board: deidentified data analysis of clinical samples was completed under Children's Mercy Hospital protocol (STUDY00000175).

1. Wojcik MH, Lemire G, Berger E, et al. Genome sequencing for diagnosing rare diseases. *N Engl J Med*. 2024;390(21):1985-1997. doi:10.1056/NEJMoa2314761

2. Cohen ASA, Farrow EG, Abdelmoity AT, et al. Genomic answers for children: dynamic analyses of >1000 pediatric rare disease genomes. *Genet Med*. 2022; 24(6):1336-1348. doi:10.1016/j.gim.2022.02.007

3. Dolzhenko E, English A, Dashnow H, et al. Characterization and visualization of tandem repeats at genome scale. *Nat Biotechnol*. 2024;42(10):1606-1614. doi:10.1038/s41587-023-02057-3

4. Chen X, Harting J, Farrow E, et al; Genomics England Research Consortium. Comprehensive SMN1 and SMN2 profiling for spinal muscular atrophy analysis using long-read PacBio HiFi sequencing. *Am J Hum Genet*. 2023;110(2):240-250. doi:10.1016/j.ajhg.2023.01.001

5. Cheung WA, Johnson AF, Rowell WJ, et al. Direct haplotype-resolved 5-base HiFi sequencing for genome-wide profiling of hypermethylation outliers in a rare disease cohort. *Nat Commun*. 2023;14(1):3090. doi:10.1038/s41467-023-38782-1

6. Groza C, Schwendinger-Schreck C, Cheung WA, et al. Pangenome graphs improve the analysis of structural variants in rare genetic diseases. *Nat Commun*. 2024;15(1):657. doi:10.1038/s41467-024-44980-2

Cannabis and Alcohol Use to Initiate Sleep Among Young Adults

More than 20% of US young adults struggle to fall or stay asleep.¹ Cannabis and alcohol can help initiate sleep,^{2,3} but regular use may be problematic.⁴ Increasing tolerance can lead to greater use to produce consistent sleep outcomes, potentially contributing to use disorder and sleep problems.²

Young adulthood is a critical developmental period for both substance use risk and sleep problems, underscoring the need for national data on cannabis and alcohol use for sleep.^{5,6}

Methods | Data were from the 2022-2023 Monitoring the Future (MTF) Panel Study collected from US young adults aged 19 to 30 years (from 2010-2022 12th-grade cohorts).⁶ MTF annually surveys nationally representative samples of 12th-grade students (modal age, 18 years); a subsample from each cohort is selected for the MTF Panel, with participation starting at age 19 or 20 years and then biennially through age 29 or 30 years.⁶ The response rate was 51.5% of active participants and 34.5% of all selected 12th graders (eFigure in Supplement 1). Written informed consent was provided by all respondents. Analysis weights accounted for sample selection and attrition. A University of Michigan Institutional Review Board approved the study. We followed the STROBE reporting guideline.

Substance use measures included past-12-month use of cannabis, alcohol, and both substances (yes or no); past 30-day daily or near-daily use (≥ 20 occasions), nondaily use (1-19 occasions), or no use; and past 2-week binge drinking (yes or no). Respondents reporting past 12-month use were asked to select all applicable answers about the primary reasons for use, including "To get to sleep."

Covariates included gender identity (man, woman, another gender), race and ethnicity (Hispanic, non-Hispanic Black [hereafter Black], non-Hispanic White [hereafter White], non-Hispanic other [Asian, Middle Eastern, multiracial, and other not specified]), age (range, 19-30 years), and college-level education (currently attending, graduated from 4-year college, other). Two-sided $P < .05$ indicated statistical significance. Data analysis was performed using SAS 9.4 (SAS Institute).

Results | Of the 1473 respondents (mean [SE] age, 24.5 [0.09] years), 51.0% identified as female, 44.7% as male, and 4.2% as another gender. Of these respondents, 11.2% identified as Black, 25.4% as Hispanic, 54.8% as White, and 8.5% as other race and ethnicity.