



OPEN Targeted metagenomics reveals hidden chickenpox epidemic amid Mpox surveillance in Uganda

Stephen Kanyerezi^{1,2,3}, Alisen Ayitewala¹, Jupiter Marina Kabahita¹, Hellen Rosette Oundo¹, Julius Sseruyange¹, Wilson Tenywa¹, Godwin Tusabe¹, Stacy Were¹, Moses Murungi¹, Martha Nabukyu^{1,2,3}, Valeria Nakintu¹, Caroline Makoha¹, Ivan Sserwadda^{2,3,5}, Harris Onywera⁴, Collins Tanui⁴, Ibrahim Mugerwa¹, Atek Kagirita⁷, Benard Lubwama⁷, Eilu Roggers Michael⁸, David Patrick Kateete³, Morgan Otita⁸, Samuel Giduddu⁹, Daudi Jjingo^{2,3}, Andrew Nsawotebba¹, Gerald Mboowa^{2,3,6}, Aloysious Ssemaganda^{1,10}, Susan Nabadda¹, Sofonias K. Tessema⁴ & Isaac Ssewanyana^{1,10}✉

In regions where both monkeypox virus (MPXV) and varicella zoster virus (VZV) are co-circulating, overlapping clinical manifestations can complicate clinical diagnosis. During the MPXV outbreak declared in Uganda in July, 2024, symptomatic suspected cases tested PCR negative for Mpox. To determine the cause of symptoms, we employed metagenomic sequencing with a targeted Viral Surveillance panel in 284 MPXV negative samples. VZV was identified as the predominant pathogen in 86% of MPXV-negative cases, suggesting a concurrent chickenpox surge. Using the VaricellaGen pipeline for variant calling, clade typing, and phylogeny, 118 (49%) samples that achieved $\geq 70\%$ genome coverage were of clade 5 based on the single-nucleotide polymorphism (SNP) dataset. This data confirms co-circulation of VZV during the MPXV outbreak in Uganda. Our results underscore the need for laboratory confirmation of MPXV and the inclusion of VZV in the testing algorithm during the Mpox outbreak.

Keywords Mpox, Varicella zoster virus (VZV), Misdiagnosis, Metagenomic sequencing, VaricellaGen, Clade 5, Uganda

Mpox is an infection caused by the monkeypox virus (MPXV), a member of the *Orthopoxvirus* genus¹. Mpox presents with a wide range of clinical symptoms such as fever, lymphadenopathy, and vesiculopustular lesions, similar to other exanthematous infections, particularly chickenpox, and has been reported in previous Mpox outbreaks^{2–5}. While MPXV can be zoonotic, VZV is predominantly human-to-human transmission with no known animal reservoir⁶. The clinical overlap between these infections makes rapid and accurate diagnosis essential for effective outbreak response and management.

VZV is a common alphaherpesvirus with a double-stranded DNA genome known to cause primarily varicella (chickenpox) in children and adolescents and herpes zoster in older and/or immunocompromised persons^{7,8}. VZV is highly contagious and is airborne with a genome of approximately 125 kb containing 71 open reading frames (ORFs)^{9,10}. Currently, nine VZV clades are recognized: seven established clades (1–6 and 9) and two provisional clades (VII and VIII)¹¹. In Mpox-endemic regions, especially in low- and middle-income countries with low chickenpox vaccination coverage, chickenpox remains commonly reported^{2–5}.

¹Department of National Health Laboratory and Diagnostics Services, Central Public Health Laboratories (NHLDS-CPHL), Ministry of Health, P.O. Box 7272, Kampala, Uganda. ²The African Center of Excellence in Bioinformatics and Data-Intensive Sciences, the Infectious Diseases Institute, College of Health Sciences, Makerere University, P.O. Box 22418, Kampala, Uganda. ³Department of Immunology and Molecular Biology, School of Biomedical Sciences, College of Health Sciences, Makerere University, P.O. Box 7072, Kampala, Uganda. ⁴Africa Centers for Disease Control and Prevention (Africa CDC), Addis Ababa, Ethiopia. ⁵Amsterdam Institute for Global Health and Development (AIGHD), Department of Global Health, Academic Medical Center, Amsterdam, The Netherlands. ⁶Broad Institute of Harvard and MIT, Cambridge, MA 02142, USA. ⁷National Public Health Emergency Operations Center-PEOC, Ministry of Health, Kampala, Uganda. ⁸Department of Global Health Security, Infectious Diseases Institute, Makerere University, Kampala, Uganda. ⁹Uganda National Institute of Public Health, Ministry of Health, Kampala, Uganda. ¹⁰Infectious Diseases Research Collaboration, Kampala, Uganda. ✉email: sewyisaac@yahoo.co.uk

Conventional laboratory methods for differentiating MPXV from VZV rely on polymerase chain reaction (PCR)-based assays, which are highly specific but limited to detecting only known targets¹². In outbreak settings, however, an overreliance on PCR may result in undetected alternative pathogens when cases test negative for the suspected virus¹³. For instance, a study during the 2014–2015 Ebola outbreak in Conakry, Guinea, found that among 2,362 suspected Ebola cases, 1,540 (65.2%) tested negative by PCR. Notably, 98 of these PCR-negative patients died, highlighting the potential presence of other severe illnesses that were initially overlooked due to the focus on Ebola¹⁴. In regions where both MPXV and VZV are co-circulating, overlapping clinical manifestations can complicate differential diagnosis, with MPXV frequently mistaken for VZV and vice versa^{2,12}. Recently, a study from Brazil deployed a multiplex PCR assay to diagnose co-infection of MPXV and VZV concurrently, but this has been limited in outbreak settings in Africa⁴. These findings underscore the need to consider alternative diagnoses and adopt comprehensive diagnostic strategies during outbreaks to avoid overlooking other clinically significant pathogens. Metagenomic sequencing offers an unbiased and powerful approach to pathogen detection, enabling the identification of viral agents beyond the initially suspected cause¹⁵.

Between September and December 2024, only 52% cases tested positive for MPXV despite presentation with symptoms. In order to understand other causes of symptoms among MPXV negatives, we utilized metagenomic sequencing.

Methods

Ethical consideration

This study was conducted in accordance with the principles outlined in the Declaration of Helsinki. All study procedures involving human participants were performed as part of the national Mpox outbreak response. Because samples were collected for public health surveillance during an active outbreak, obtaining individual informed consent was not feasible. Consequently, a waiver of informed consent was sought and approved by the Uganda National Health Laboratory Services Research and Ethics Committee (UNHLS-2025-133).

Sample collection and PCR testing

During the MPXV outbreak declared in Uganda in July, 2024, 284 symptomatic suspected cases tested PCR negative for Mpox between September and December 2024 were included in this study. Suspected Mpox cases were identified in accordance with the World Health Organization (WHO) outbreak case definition. Briefly, suspected cases included individuals presenting with an unexplained acute skin rash, mucosal lesions, and/or lymphadenopathy, with or without systemic symptoms such as fever, headache, myalgia, back pain, or profound weakness, or individuals with epidemiological linkage to a probable or confirmed Mpox case within 21 days prior to symptom onset. Consistent with WHO guidance, common causes of acute rash illness such as varicella (chickenpox), herpes zoster, and measles were considered during clinical assessment but did not preclude MPXV testing, given documented clinical overlap and the possibility of co-infections¹⁶. The samples included lesion, rash, oropharyngeal, or genital swabs in Viral Transport Media (VTM), as well as crusts collected in plain tubes - obtained as part of the public health emergency response to Mpox from affected individuals across 27 districts in the country. DNA was extracted using QIAamp DNA Kit (Qiagen, Hilden, Germany) and subsequently subjected to the Non-Variola Orthopoxvirus Real-Time PCR Primer and Probe Set (CDC, EUA)¹⁷ in-house assay at the Central Emergency Response and Surveillance Laboratory (CERSL), Ministry of Health, Uganda. Briefly, amplification was conducted on the CFX96 Connect Real-Time Detection System (Bio-Rad Laboratories, Inc., California, USA), with denaturation at 95 °C for 8 min, followed by 40 cycles, each consisting of two main steps i.e., a denaturation step at 95 °C for 5 s and an annealing and extension step at 60 °C for 30 s. PCR results were analyzed using CFX96 Real-Time PCR system (Bio-Rad Laboratories, Inc., California, USA) for Ct value interpretation and quality control.

Metagenomic sequencing and taxonomic classification

High-quality DNA was extracted using the QIAamp DNA Kit (Qiagen, Hilden, Germany), normalized to a starting concentration of 50 ng, and prepared for metagenomic sequencing using a target enrichment approach with a pre-designed Illumina Viral Surveillance Panel (VSP). Paired-end sequencing was performed on the Illumina MiSeq platform at the Genomics Core Laboratory of the Central Public Health Laboratory. Raw sequencing reads were analyzed using KrakenUniq (v1.0.3) for taxonomic classification and identification of viral pathogens¹⁸. Using the developed VaricellaGen pipeline, reads were checked for quality, consensus genomes generated and only samples that attained genome coverage of 70% and above were subjected to downstream analysis. The same pipeline was used to assign clades to genomes and construct a phylogenetic tree. The resulting phylogenetic tree was visualized using the Interactive Tree of Life (iTOL v7) platform for intuitive exploration and annotation of clade relationships¹⁹.

VaricellaGen: automated VZV genomic analysis pipeline development

To enhance the genomic characterization of VZV, we developed VaricellaGen (<https://github.com/MicroBioGenoHub/VaricellaGen>). The pipeline includes modules for quality control, variant calling, consensus genome generation, clade typing using a curated SNP dataset¹¹, and phylogenetic analysis with 1000 bootstraps and collapsing near to zero branch lengths. For the phylogenetic module, reference VZV sequences with known clade assignments were retrieved from NCBI and concatenated into a single FASTA file to serve as a background dataset^{11, 20}–³². The sequence JN704693.1 was designated as the outgroup. Sequence identifiers were standardized and annotated in the following format: GenBank accession, three-letter country code of sample origin, year of collection, and assigned clade.

Variant calling and consensus genome assembly

The quality assessment of sequencing reads was designed to use FastQC (v0.12.1) and poor quality reads trimmed using Trim Galore (v0.6.10). The alignment of passed reads were aligned to the VZV reference genome (NC_001348.1) using bwa (v0.7.18-r1243-dirty) and samtools (v1.21), and variants are called using freebayes (v1.3.8) and BCFtools (v1.21)^{33–37}. Process_gvcf from ARTIC was used to generate regions to be masked during consensus genome generation³⁸. To generate consensus sequences, VaricellaGen was designed to use BCFtools, and only samples that attained genome coverage of 70% and above were meant to be subjected to downstream analysis.

Clade assignment and phylogenetic analysis

To classify the VZV strains, VaricellaGen used an integrated SNP-based clade typing module. For phylogenetic analysis, consensus genomes are concatenated and processed through VaricellaGen's phylogeny module, which performs multiple sequence alignment using MAFFT (v7.525) and constructs a maximum-likelihood phylogenetic tree with IQ-TREE2 (v2.4.0) to assess the genetic relatedness of query samples to previously reported VZV strains^{39,40}. VaricellaGen uses reference VZV sequences with known clade assignments retrieved from NCBI and concatenated into a single FASTA file to serve as a background dataset. The sequence JN704693.1 was designated as the outgroup. Sequence identifiers were standardized and annotated in the following format: GenBank accession, three-letter country code of sample origin, year of collection, and assigned clade.

Results

In this study, we analyzed 284 PCR-confirmed MPXV-negative cases from 27 districts (Fig. 1). These included 151 (53.2%) males and 123 (43.3%) females, with a mean age of 14 years (range: 9 months–70 years), who




Fig. 1. Geographic distribution of sample collection sites across Uganda. Map of Uganda showing district-level Mpox PCR positivity rates based on national surveillance data collected between July and December 2024, alongside the 27 districts from which MPXV-negative clinical samples were collected for metagenomic sequencing. The map was generated using QGIS (version QGIS-OSGeo4W-3.44.6-1; <https://qgis.org/>). Districts with no reported data or no samples tested during the study period are not labeled.

presented with Mpox-like symptoms, notably vesicular-pustular rash, fever, and lymphadenopathy (<https://bit.ly/4iZeQDn>). Gender information was unavailable for 10 (3.5%) isolates (Table 1).

PCR screening ruled out MPXV infection in all clinically suspected cases. Metagenomic sequencing, followed by taxonomic classification using KrakenUniq, identified VZV as the predominant viral pathogen. Specifically, 86% (243/284) of the samples had at least one read aligning to VZV, while 14% (41/284) showed no VZV reads. No other clinically significant DNA viruses were consistently detected among MPXV and VZV negative samples within the limits of the DNA-based metagenomic approach used.

VaricellaGen: automated VZV genomic analysis pipeline

To enhance the genomic characterization of VZV, we successfully developed VaricellaGen (<https://github.com/MicroBioGenoHub/VaricellaGen>), an automated pipeline for the genomic analysis of VZV. The pipeline integrates sequential modules for raw data quality control, variant calling, consensus genome generation, SNP-based clade typing¹¹, and phylogenetic reconstruction with 1000 bootstraps and collapsing near to zero branch lengths (Fig. 2). Outputs include annotated consensus genomes, clade assignments, and a phylogenetic tree, providing a streamlined workflow from raw reads to interpretable genomic insights.

Clade assignment and phylogenetic insights

Of the 243 VZV-positive samples, 49% (118/243) achieved ≥70% genome coverage using the VaricellaGen pipeline. SNP-based clade typing classified all successfully sequenced samples as VZV clade 5. This finding was corroborated by phylogenetic analysis, which revealed two distinct sub-clusters within clade 5: one cluster predominantly composed of sequences from Ugandan, India, United States, and Kenyan samples; and a second cluster comprising the remaining Ugandan sequences along with one sequence from Indian (Fig. 3).

Discussion

Genomic pathogen surveillance is critically important, especially during outbreaks with non-specific or overlapping clinical presentations. Here, we investigated cases that presented with Mpox-like signs but had MPXV infection ruled out by PCR. Metagenomic sequencing revealed VZV as the predominant etiological agent in a substantial proportion of cases, suggesting a concurrent and previously unrecognized surge in chickenpox infections during the Mpox outbreak in Uganda.

These findings are consistent with concurrent MPXV and VZV outbreaks reported in Burundi, Kenya, and the DRC^{41–43} and emphasize the challenge of differentiating co-circulating rash-causing viruses based on clinical signs alone^{44,45}. The similar skin manifestations of MPXV and VZV can complicate clinical diagnosis, especially in resource-limited settings with limited access to specific tests. This overlap underscores the necessity of comprehensive laboratory diagnostics, beyond single PCR assays, for accurate identification of the causative agent and effective public health responses during outbreaks.

Using a metagenomic sequencing approach, we identified VZV in 86% of PCR-confirmed MPXV-negative cases that were clinically suspected to be Mpox. Our newly developed VaricellaGen pipeline enabled detailed genomic analysis, showing that all samples with sufficient genome coverage belonged to VZV clade 5. Phylogenetic clustering with established clade 5 reference genomes further supported this classification and suggests that this clade is prevalent in Uganda, consistent with previous findings from other African countries^{32,46}.

Our phylogenetic analysis of VZV clade 5 revealed two Ugandan sub-clusters. One mainly contained sequences from Uganda, India, US, and Kenya, suggesting regional lineage mixing or historical introductions. The other primarily included the remaining Ugandan sequences, clustering with one isolate from India, indicative of multiple introductions or diverse transmission chains within Uganda. This clustering pattern also points to travel-related spread or shared networks across distant regions. These findings highlight the importance of detailed genomic surveillance for both local and international transmission tracking. The observed sub-structuring within clade 5 strongly suggests the need for a standard naming system for VZV sub-clusters to improve genomic reporting and the precision of transmission analysis for future outbreak management.

The simultaneous detection of VZV and MPXV outbreaks highlights critical gaps in current outbreak response protocols, particularly in differential diagnosis and surveillance systems. Large-scale epidemics often focus on a single high-priority pathogen, inadvertently leading to missing concurrent infections, a pattern previously seen in influenza pandemics with underrecognized bacterial co-infections despite their significant

Variable	Value
Total cases analyzed	284
MPXV status	PCR-confirmed MPXV-negative
Number of districts	27
Sex distribution	151 (53.2%) males and 123 (43.3%) females. Gender information unavailable for 10 cases (3.5%)
Mean age	14 years
Age range	9 months – 70 years
Clinical presentation	Vesicular-pustular rash, fever, lymphadenopathy

Table 1. Demographic and clinical characteristics of 284 PCR-confirmed MPXV-negative cases presenting with Mpox-like symptoms.




Fig. 2. Flowchart of the VaricellaGen pipeline. The pipeline integrates sequential modules for raw data quality control, variant calling, consensus genome generation, clade typing (based on¹¹, and phylogenetic reconstruction using a curated background dataset. Outputs include clade assignments, annotated consensus genomes, and phylogenetic trees for downstream interpretation.

contributions to morbidity and mortality⁴⁷. These findings emphasize the need for integrated surveillance strategies with the ability to capture a wide spectrum of pathogens underscoring the need for multi-pathogen molecular diagnostics and genomic tools to avoid misclassification, especially in sub-Saharan Africa, where integrated disease surveillance systems have previously demonstrated the limitations of single-pathogen monitoring⁴⁸. A One Health approach, at the human, animal, and environment intersection further emphasizes the importance of monitoring multiple pathogens to detect emerging and re-emerging threats⁴⁹. Our findings support the growing call for unbiased sequencing and comprehensive diagnostics in both routine surveillance and emergency responses.

Despite VZV being a DNA virus with a generally slow mutation rate⁵⁰, our analyzed sequences showed moderate genetic diversity (Fig. 3), suggesting ongoing viral evolution or multiple introduction events. Since all sequences belonged to clade 5, the observed variation could indicate a within-clade origin reflective of probable local transmission patterns, accumulated mutations over time, or different points of origin.

Strengths of this study lie in the targeted DNA-based metagenomic sequencing, which enabled comprehensive detection and genomic characterization of DNA viruses, particularly VZV, among MPXV-negative cases, and the development of VaricellaGen, an automated pipeline for the rapid genomic characterization of VZV. By integrating quality control, variant calling, clade typing, and phylogenetics, VaricellaGen offers a scalable tool for outbreak genomic epidemiology. A key limitation of our study was that detailed genomic analyses only included samples with $\geq 70\%$ genome coverage. While this threshold ensured data quality, it may have excluded informative lower-quality samples. In addition, the use of DNA-only extraction for metagenomic sequencing restricted pathogen detection to DNA viruses and excluded RNA viruses such as enteroviruses and alphaviruses (e.g., Sindbis virus), several of which are well-recognized causes of vesicular rash illness, underscoring the importance of robust sample collection, handling, and comprehensive nucleic acid extraction strategies⁵¹. Future investigations incorporating total nucleic acid extraction will be required to comprehensively evaluate alternative viral etiologies of vesicular rash illness, while further research should also explore whether specific genomic features within VZV clade 5 are associated with disease severity, transmission dynamics, or geographic spread.

Overall, our findings emphasize the crucial role of integrated genomic surveillance during outbreaks, especially when clinical presentation cannot be relied on to differentiate between circulating pathogens. The identification of VZV during Uganda's Mpox outbreak demonstrates the value of metagenomic sequencing approaches for investigating alternative causes of Mpox-like illness and emphasizes the importance of strengthening diagnostic




Fig. 3. Phylogenetic analysis of Varicella Zoster Virus (VZV) genomes from Uganda. **(A)** Maximum-likelihood phylogenetic tree of VZV genomes. Phylogenetic analysis of VZV-positive samples from Uganda (green) alongside reference sequences from GenBank representing known clades (Clades 1–9). **(B)** All Ugandan samples (in red) clustered within Clade 5, forming two distinct sub-clusters: one grouping closely with sequences from India, the United States, and Kenya, and another clustering with only one sequence from India. The tree was constructed using IQ-TREE2 with MAFFT-based alignments; JN704693.1 (Clade 1) was used as the outgroup. Bootstrap values > 30 are shown at key nodes. Tree visualization was performed using iTOL.

capacity during outbreaks. This study further illustrates how genomic tools can aid in differential diagnosis, guide public health response, and reduce the risks of diagnostic ambiguity. The development of VaricellaGen offers a scalable, automated tool for detailed VZV genomic surveillance, providing a valuable resource for future outbreak investigations and public health decision-making.

Data availability

The source code and operation manual for VaricellaGen are available from GitHub under GNU GPL v3; (<https://github.com/MicroBioGenoHub/VaricellaGen>). The authors confirm that all supporting data, code, and protocols have been provided within the article. The genomic raw read files from this study are publicly available at the Sequence Read Archive (SRA) of the National Center for Biotechnology Information (NCBI) under the study BioProject ID: PRJNA1373217.

Received: 9 December 2025; Accepted: 30 January 2026

Published online: 06 February 2026

References

- Shchelkunova, G. A. & Shchelkunov, S. N. Smallpox, Monkeypox and other human orthopoxvirus infections. *Viruses* **15** (1), 103. <https://doi.org/10.3390/v15010103> (Dec. 2022).
- Hoff, N. A. et al. Varicella coinfection in patients with active Monkeypox in the Democratic Republic of the Congo. *EcoHealth* **14** (3), 564–574. <https://doi.org/10.1007/s10393-017-1266-5> (Sept. 2017).
- Hughes, C. M. et al. Feb., A Tale of Two Viruses: Coinfections of Monkeypox and Varicella Zoster Virus in the Democratic Republic of Congo. *Am. J. Trop. Med. Hyg.*, vol. 104, no. 2, pp. 604–611, (2021). <https://doi.org/10.4269/ajtmh.20-0589>
- Martins-Filho, P. R. et al. First reports of Monkeypox and varicella-zoster virus coinfection in the global human Monkeypox outbreak in 2022. *Travel Med. Infect. Dis.* **51**, 102510. <https://doi.org/10.1016/j.tmaid.2022.102510> (2023).
- Tayachew, A. et al. Genomic evidence of varicella-zoster virus among Mpox-suspected cases in Ethiopia during the 2022 Mpox multi-country outbreak. *Sci. Rep.* <https://doi.org/10.1038/s41598-025-29116-w> (Nov. 2025).
- Di Giulio, D. B. & Eckburg, P. B. Human monkeypox: an emerging zoonosis. *Lancet Infect. Dis.* **4** (1), 15–25. [https://doi.org/10.1016/S1473-3099\(03\)00856-9](https://doi.org/10.1016/S1473-3099(03)00856-9) (Jan. 2004).
- Gershon, A. A. et al. Varicella Zoster virus infection. *Nat. Rev. Dis. Primer.* **1**, 15016. <https://doi.org/10.1038/nrdp.2015.16> (July 2015).
- Wharton, M. The epidemiology of varicella-zoster virus infections. *Infect. Dis. Clin. North. Am.* **10** (3), 571–581. [https://doi.org/10.1016/s0891-5520\(05\)70313-5](https://doi.org/10.1016/s0891-5520(05)70313-5) (Sept. 1996).
- Gershon, A. A. Varicella zoster vaccines and their implications for development of HSV vaccines, *Virology*, vol. 435, no. 1, pp. 29–36, Jan. (2013). <https://doi.org/10.1016/j.virol.2012.10.006>
- Breuer, J. Molecular Genetic Insights Into Varicella Zoster Virus (VZV), the vOka Vaccine Strain, and the Pathogenesis of Latency and Reactivation, *J. Infect. Dis.*, vol. 218, no. suppl_2, pp. S75–S80, Sept. (2018). <https://doi.org/10.1093/infdis/jiy279>
- Jensen, N. J. et al. Revisiting the genotyping scheme for varicella-zoster viruses based on whole-genome comparisons. *J. Gen. Virol.* **98**, 1434–1438. <https://doi.org/10.1099/jgv.0.000772> (June 2017).
- Balingit, A. K. et al. Nov., Clinical presentation and molecular diagnosis of a possible Mpox virus and Varicella zoster virus coinfection in an adult immunocompetent Filipino: a case report, *Front. Public Health*, vol. 12, p. 1387636, (2024). <https://doi.org/10.3389/fpubh.2024.1387636>
- Mortier, C. et al. How to distinguish Mpox from its mimickers: an observational retrospective cohort study. *J. Med. Virol.* **95** (10), e29147. <https://doi.org/10.1002/jmv.29147> (2023).
- Ingelbeen, B. et al. Mortality among PCR negative admitted Ebola suspects during the 2014/15 outbreak in Conakry, Guinea: A retrospective cohort study. *PLoS ONE* **12** (6), e0180070. <https://doi.org/10.1371/journal.pone.0180070> (June 2017).
- Gu, W., Miller, S. & Chiu, C. Y. Clinical metagenomic Next-Generation sequencing for pathogen detection. *Annu. Rev. Pathol.* **14**, 319–338. <https://doi.org/10.1146/annurev-pathmechdis-012418-012751> (Jan. 2019).
- WHO case definitions for mpox outbreak in non-endemic countries, in Clinical management and infection prevention and control for mpox: living guideline [Internet], World Health Organization. Accessed: Jan. 29, 2026. [Online]. (2025). Available: <https://www.ncbi.nlm.nih.gov/books/NBK616033/>
- CDC. Non-variola Orthopoxvirus Real-time PCR Primer and Probe Set - EUA. (2024).
- Breitwieser, F. P., Baker, D. N. & Salzberg, S. L. KrakenUniq: confident and fast metagenomics classification using unique k-mer counts. *Genome Biol.* **19** (1), 198. <https://doi.org/10.1186/s13059-018-1568-0> (Nov. 2018).
- Leticia, I. & Bork, P. Interactive Tree Of Life (iTOL): an online tool for phylogenetic tree display and annotation, *Bioinforma. Oxf. Engl.*, vol. 23, no. 1, pp. 127–128, Jan. (2007). <https://doi.org/10.1093/bioinformatics/btl529>
- Human alphaherpesvirus 3 isolate KE_23082024, complete genome. Nov. 24. Accessed: Apr. 15, 2025. [Online]. (2024). Available: <http://www.ncbi.nlm.nih.gov/nuccore/PQ505474.1>
- Gomi, Y. et al. Comparison of the complete DNA sequences of the Oka varicella vaccine and its parental virus, *J. Virol.*, vol. 76, no. 22, pp. 11447–11459, Nov. (2002). <https://doi.org/10.1128/jvi.76.22.11447-11459.2002>
- Santos, R. A., Padilla, J. A., Hatfield, C. & Grose, C. Antigenic variation of varicella zoster virus Fc receptor gE: loss of a major B cell epitope in the ectodomain, *Virology*, vol. 249, no. 1, pp. 21–31, Sept. (1998). <https://doi.org/10.1006/viro.1998.9313>
- Complete-Genome Phylogenetic Approach to Varicella-Zoster Virus Evolution. Genetic Divergence and Evidence for Recombination, CoLab. Accessed: Apr. 15, 2025. [Online]. Available: <https://colab.ws/articles/10.1128%2Fjvi.00835-06>
- Peters, G. A. et al. Oct., A full-genome phylogenetic analysis of varicella-zoster virus reveals a novel origin of replication-based genotyping scheme and evidence of recombination between major circulating clades, *J. Virol.*, vol. 80, no. 19, pp. 9850–9860, (2006). <https://doi.org/10.1128/JVI.00715-06>
- Human herpesvirus 3 strain VZV/Philadelphia.USA/73/V[3], partial genome. May 16, 2012. Accessed: Apr. 15, 2025. [Online]. Available: <http://www.ncbi.nlm.nih.gov/nuccore/JQ911599.1>
- Garcés-Ayala, F. et al. Full-Genome sequence of a novel Varicella-Zoster virus clade isolated in Mexico. *Genome Announc.* **3**, 4, pp. e00752–15, July 2015. <https://doi.org/10.1128/genomeA.00752-15>
- Depledge, D. P. et al. Evolution of cocirculating Varicella-Zoster virus genotypes during a chickenpox outbreak in Guinea-Bissau. *J. Virol.* **88**, 13936–13946. <https://doi.org/10.1128/JVI.02337-14> (Dec. 2014).
- Norberg, P. et al. Recombination of globally Circulating Varicella-Zoster virus. *J. Virol.* **89** (14), 7133–7146. <https://doi.org/10.1128/JVI.00437-15> (Apr. 2015).
- Jensen, N. J. et al. Analysis of the reiteration regions (R1 to R5) of varicella-zoster virus. *Virology* **546**, 38–50. <https://doi.org/10.1016/j.virol.2020.03.008> (July 2020).
- Davison, A. J. & Scott, J. E. The complete DNA sequence of varicella-zoster virus. *J. Gen. Virol.* **67**, 1759–1816. <https://doi.org/10.1099/0022-1317-67-9-1759> (Sept. 1986). Pt 9.

31. El-Duah, P. et al. Genetic characterization of varicella-zoster and HIV-1 viruses from the cerebrospinal fluid of a co-infected encephalitic patient, Ghana. *Virol. J.* **19**, 122. <https://doi.org/10.1186/s12985-022-01854-7> (July 2022).
32. Kumar, A. et al. First detection of varicella Zoster virus clade 9 cases in India during Mpox surveillance. *Ann Med*, **55**, 2, p. 2253733, <https://doi.org/10.1080/07853890.2023.2253733>
33. Danecek, P. et al. Feb., Twelve years of SAMtools and BCFtools, *GigaScience*, vol. 10, no. 2, p. giab008, (2021). <https://doi.org/10.1093/gigascience/giab008>
34. Garrison, E. & Marth, G. Haplotype-based variant detection from short-read sequencing, *ArXiv12073907 Q-Bio*, July Accessed: Mar. 25, 2022. [Online]. (2012). Available: <http://arxiv.org/abs/1207.3907>
35. Krueger, F. *FelixKrueger/TrimGalore* Feb. 02., Perl. Accessed: Feb. 03, 2025. [Online]. (2025). Available: <https://github.com/FelixKrueger/TrimGalore>
36. LaMar, D. & FastQC doi: (2015). <https://qubeshub.org/resources/fastqc>
37. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows–Wheeler transform. *Bioinformatics* **25**, 1754–1760. <https://doi.org/10.1093/bioinformatics/btp324> (July 2009).
38. BCCDC-PHL BCCDC-PHL/mpxv-artic-nf. Accessed: Apr. 08, 2025. [Online]. Available: <https://github.com/BCCDC-PHL/mpxv-artic-nf>
39. Minh, B. Q. et al. IQ-TREE 2: new models and efficient methods for phylogenetic inference in the genomic era. *Mol. Biol. Evol.* **37** (5), 1530–1534. <https://doi.org/10.1093/molbev/msaa015> (May 2020).
40. Katoh, K. & Standley, D. M. MAFFT Multiple Sequence Alignment Software Version 7: Improvements in Performance and Usability, *Mol. Biol. Evol.*, vol. 30, no. 4, pp. 772–780, Apr. (2013). <https://doi.org/10.1093/molbev/mst010>
41. Leung, J. et al. Varicella in the Tshuapa Province, Democratic Republic of Congo, 2009–2014. *Trop. Med. Int. Health TM IH*. **24** (7), 839–848. <https://doi.org/10.1111/tmi.13243> (July 2019).
42. Onyango, C. et al. High prevalence of varicella zoster virus infection among persons with suspect mpox cases during a mpox outbreak in Kenya, 2024, Apr. 10, *medRxiv*. (2025). <https://doi.org/10.1101/2025.04.08.25325502>
43. Nzoyikorera, N. et al. Monkeypox clade Ib virus introduction into burundi: first findings, July to mid-August 2024. *Eurosurveillance* **29** (42), 2400666. <https://doi.org/10.2807/1560-7917.ES.2024.29.42.2400666> (Oct. 2024).
44. Derrick, K. M., Marson, J. W., Chakka, S. & Heilman, E. R. Mpox virus and coinfections: An approach to rapid diagnosis, *J. Cutan. Pathol.*, vol. 50, no. 10, pp. 878–883, Oct. (2023). <https://doi.org/10.1111/cup.14490>
45. Guillén-Calvo, L. et al. Mpox, herpes, and enteroviruses: differential diagnosis. *J. Med. Virol.* **96** (1), e29371. <https://doi.org/10.1002/jmv.29371> (2024).
46. Bryant, P. et al. Vaccine strain and Wild-Type clades of Varicella-Zoster virus in central nervous system and Non-CNS Disease, new York State, 2004–2019. *J. Clin. Microbiol.*, **60**, 4, pp. e02381–e02321, <https://doi.org/10.1128/jcm.02381-21>
47. Morris, D. E., Cleary, D. W. & Clarke, S. C. Secondary bacterial infections associated with influenza pandemics. *Front. Microbiol.* **8**, 1041. <https://doi.org/10.3389/fmicb.2017.01041> (2017).
48. Mremi, I. R. et al. Twenty years of integrated disease surveillance and response in Sub-Saharan africa: challenges and opportunities for effective management of infectious disease epidemics. *One Health Outlook*. **3** (1), 22. <https://doi.org/10.1186/s42522-021-00052-9> (Nov. 2021).
49. Kelly, T. R. et al. Implementing one health approaches to confront emerging and re-emerging zoonotic disease threats: lessons from PREDICT. *One Health Outlook*. **2** (1, p. 1,). <https://doi.org/10.1186/s42522-019-0007-9> (Jan. 2020).
50. Peters, G. A. et al. Oct., The Attenuated Genotype of Varicella-Zoster Virus Includes an ORF0 Transitional Stop Codon Mutation, *J. Virol.*, vol. 86, no. 19, pp. 10695–10703, (2012). <https://doi.org/10.1128/jvi.01067-12>
51. Ajay, S. S., Parker, S. C. J., Ozel Abaan, H., Fuentes Fajardo, K. V. & Margulies, E. H. Accurate and comprehensive sequencing of personal genomes, *Genome Res.*, vol. 21, no. 9, pp. 1498–1505, Sept. (2011). <https://doi.org/10.1101/gr.123638.111>

Acknowledgements

We acknowledge the Ministry of Health Uganda and the Central Public Health Laboratories (CPHL) for supporting sample collection, diagnostic testing, and genomic sequencing during the Mpox outbreak response. We are grateful to the Genomics Core Facility at CPHL for sequencing support and to the field epidemiology and surveillance teams for their commitment to timely case investigation and sample coordination. We also thank the Africa Centers for Disease Control and Prevention (Africa CDC) for their technical support and continued investment in strengthening genomic surveillance capacity in Uganda.

Author contributions

Conceptualization: S.K., A.A., I.S., S.N., D.J., G.M., D.P.K., A.S., S.K.T., I.S. Formal analysis: S.K., A.A., J.M.K., C.M., I.S., G.M. Funding acquisition: S.N., I.S., S.T.K., H.O., C.T., N.A., A.A., A.S., S.K., I.M., A.K. Methodology: S.K., A.A., J.M.K., H.R.O., J.S., W.T., G.T., S.W., M.M., M.N., V.Z.N., C.M., I.S., G.M., A.S., I.M., B.L., E.M.R., M.O., S.G. Validation: S.K., A.A., B.A.K., H.O., C.T., N.A., V.Z.N., N.A., G.M., A.S., S.K.T., I.S. Visualization: S.K., I.S., J.M.K. Software: S.K., J.M.K., I.S., G.M. Writing – original draft: S.K., A.A., B.A.K., V.Z.N., I.S., H.O., C.T., N.A., S.N., D.J., G.M., A.A., S.K.T., I.S. Writing – review and editing: S.K., A.A., B.A.K., V.Z.N., I.S., H.O., C.T., N.A., S.N., D.J., G.M., A.A., S.K.T., I.S., I.M., B.L., E.M.R., M.O., S.G., A.K., D.P.K.

Funding

This work was supported by public health emergency response funds from the Government of Uganda.

Declarations

Competing interests

The authors declare no competing interests.

Additional information

Correspondence and requests for materials should be addressed to I.S.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Open Access This article is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License, which permits any non-commercial use, sharing, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if you modified the licensed material. You do not have permission under this licence to share adapted material derived from this article or parts of it. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by-nc-nd/4.0/>.

© The Author(s) 2026