



DATA NOTE

REVISED The genome sequence of the Antarctic lanternfish,

Electrona antarctica (Günther, 1878)

[version 2; peer review: 3 approved]

Iliana Bista ¹, Martin Collins ²,

Wellcome Sanger Institute Tree of Life Management, Samples and Laboratory team,

Wellcome Sanger Institute Scientific Operations: Sequencing Operations,
Wellcome Sanger Institute Tree of Life Core Informatics team,
Tree of Life Core Informatics collective¹Senckenberg Research Institute and Natural History Museum, Frankfurt, Germany²British Antarctic Survey, NERC, Cambridge, England, UK
v2 First published: 19 Feb 2025, 10:89
<https://doi.org/10.12688/wellcomeopenres.23803.1>

 Latest published: 25 Jul 2025, 10:89
<https://doi.org/10.12688/wellcomeopenres.23803.2>
Abstract

We present a genome assembly from an individual female *Electrona antarctica* (the Antarctic lanternfish; Chordata; Actinopterygii; Myctophiformes; Myctophidae). The genome sequence has a total length of 1,427.40 megabases. Most of the assembly is scaffolded into 24 chromosomal pseudomolecules. The mitochondrial genome has also been assembled and is 20.02 kilobases in length.

Keywords
Electrona antarctica, Antarctic lanternfish, genome sequence, chromosomal, Myctophiformes

 This article is included in the [Tree of Life gateway](#).
Open Peer Review**Approval Status**

	1	2	3
version 2 (revision) 25 Jul 2025			 view
version 1 19 Feb 2025	 view	 view	 view

 1. **Takushi Kishida** , Nihon University,
Fujisawa, Japan

 2. **Jayan Duminda M. Senevirathna** , Uva
Wellassa University, Badulla, Sri Lanka

 3. **Annabel Whibley** , Bragato Research
Institute, Lincoln, New Zealand

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding author: Wellcome Sanger Institute Tree of Life Management, Samples and Laboratory team (Mark.Blaxter@sanger.ac.uk)

Author roles: **Bista I:** Investigation, Resources, Writing – Original Draft Preparation, Writing – Review & Editing; **Collins M:** Resources, Writing – Review & Editing;

Competing interests: No competing interests were disclosed.

Grant information: This work was supported by Wellcome through core funding to the Wellcome Sanger Institute (220540). Specimen collection was supported by a Collaborative Antarctic Science Scheme fund (CASS) (CASS-174/2019) by the British Antarctic Survey awarded to IB. IB was supported by the Centre for Translational Biodiversity Genomics (LOEWE-TBG) through the program LOEWE–Landes-Offensive zur Entwicklung Wissenschaftlich-ökonomischer Exzellenz of Hesse's Ministry of Higher Education, Research, and the Arts.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2025 Bista I *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Bista I, Collins M, Wellcome Sanger Institute Tree of Life Management, Samples and Laboratory team *et al.* **The genome sequence of the Antarctic lanternfish, *Electrona antarctica* (Günther, 1878) [version 2; peer review: 3 approved]** Wellcome Open Research 2025, 10:89 <https://doi.org/10.12688/wellcomeopenres.23803.2>

First published: 19 Feb 2025, 10:89 <https://doi.org/10.12688/wellcomeopenres.23803.1>

REVISED Amendments from Version 1

The text describing the PacBio read coverage number of reads and number of Gigabases has been corrected to include all of the reads.

Any further responses from the reviewers can be found at the end of the article

Species taxonomy

Eukaryota; Opisthokonta; Metazoa; Eumetazoa; Bilateria; Deuterostomia; Chordata; Craniata; Vertebrata; Gnathostomata; Teleostomi; Euteleostomi; Actinopterygii; Actinopteri; Neopterygii; Teleostei; Osteoglossocephalai; Clupeocephala; Euteleostei; Neoteleostei; Eurypterygia; Ctenosquamata; Myctophata; Myctophiformes; Myctophidae; *Electrona*; *Electrona antarctica* (Günther, 1878) (NCBI:txid206093).

Background

The Antarctic lanternfish *Electrona antarctica*, is a member of the Myctophidae family of fish (Order: Myctophiformes). Myctophidae are mesopelagic fish comprising over 250 species and constitute one of the most dominant fish groups throughout the global oceans (Catul *et al.*, 2011; Collins *et al.*, 2012). Myctophids undergo extensive vertical migration moving between depths of 50 and 400 m during different times of the day (Hulley & Duhamel, 2011). Feeding in myctophids is aimed mostly at mesozooplankton, including amphipods, copepods, and euphasids (krill), while they are themselves major feeding targets of penguins, other fish (e.g. toothfish), seals and other higher predators (Collins *et al.*, 2012).

Populations of the Antarctic lanternfish *E. antarctica* occur in the Southern Ocean (SO), with a distribution including the Antarctic Polar front and extending all the way to the far South (Hulley, 1990). *E. antarctica* is considered a key species, among the already dominant Myctophidae family, due to its high abundance and wide distribution. The conservation status of this species is currently listed as a “Least concern” in the IUCN red list (Williams, 2019).

Here we present a genome assembly generated using an adult specimen collected from the Antarctic Polar Front area, during the JR19001 cruise onboard the James Clark Ross research vessel.

Genome sequence report

The genome of a specimen of *Electrona antarctica* (Figure 1) was sequenced using Pacific Biosciences single-molecule HiFi long reads, generating a total of 54.73 Gb (gigabases) from 6.17 million reads, providing approximately 43-fold coverage. Primary assembly contigs were scaffolded with chromosome conformation Hi-C data, which produced 251.34 Gb from 1,664.50 million reads, yielding an approximate coverage of 176-fold. Specimen and sequencing details are provided in Table 1.



Figure 1. Photograph of the *Electrona antarctica* (fEleAnt2) specimen used for genome sequencing.

Manual assembly curation corrected 100 missing joins or mis-joins and 12 haplotypic duplications, reducing the scaffold number by 2.11%, and increasing the scaffold N50 by 2.99%. The final assembly has a total length of 1,427.40 Mb in 1,900 sequence scaffolds with a scaffold N50 of 53.6 Mb (Table 2). The total count of gaps in the scaffolds is 3,179. The snail plot in Figure 2 provides a summary of the assembly statistics, while the distribution of assembly scaffolds on GC proportion and coverage is shown in Figure 3. The cumulative assembly plot in Figure 4 shows curves for subsets of scaffolds assigned to different phyla. Most (88.72%) of the assembly sequence was assigned to 24 chromosomal-level scaffolds. Chromosome-scale scaffolds confirmed by the Hi-C data are named in order of size (Figure 5; Table 3). While not fully phased, the assembly deposited is of one haplotype. Contigs corresponding to the second haplotype have also been deposited. The mitochondrial genome was also assembled and can be found as a contig within the multifasta file of the genome submission.

The estimated Quality Value (QV) of the final assembly is 53.3. The *k*-mer completeness was estimated at 74.95% for the primary assembly, 71.26% for the alternate haplotype and 96.62% for the combined assemblies. The primary assembly achieves a BUSCO v5.3.2 completeness of 95.9% (single = 83.1%, duplicated = 12.7%), using the actinopterygii_odb10 reference set (*n* = 3,640).

Methods

Sample acquisition

Specimens of *Electrona antarctica* were collected from the Southern Ocean (latitude −53.36, longitude −38.06) on 2019-12-05. The specimen was collected by Iliana Bista and identified by Martin Collins (British Antarctic Survey) using a deep-sea net. Individual tissues were dissected from the euthanised specimen and preserved immediately in the a −80°C freezer until further processing.

The genome sequence is based on PacBio HiFi sequencing data from specimen ID SAN1100012 (ToLID fEleAnt2). A second specimen (specimen ID SAN1100013, ToLID fEleAnt3) was used for Hi-C and RNA sequencing.

Table 1. Specimen and sequencing data for *Electrona antarctica*.

Project information			
Study title	<i>Electrona antarctica</i> (Antarctic lanternfish)		
Umbrella BioProject	PRJEB60649		
Species	<i>Electrona antarctica</i>		
BioSample	SAMEA8748807		
NCBI taxonomy ID	206093		
Specimen information			
Technology	ToLID	BioSample accession	
PacBio long read sequencing	fEleAnt2	SAMEA8748807	
Hi-C sequencing	fEleAnt3	SAMEA8748808	
RNA sequencing	fEleAnt3	SAMEA8748826	
Sequencing information			
Platform	Run accession	Read count	Base count (Gb)
Hi-C Illumina NovaSeq 6000	ERR11040177	1.66e+09	251.34
PacBio Sequel IIe	ERR11029670	2.18e+06	19.29
PacBio Sequel IIe	ERR11029669	9.39e+05	10.84
PacBio Sequel IIe	ERR11029671	1.47e+06	10.94
PacBio Sequel IIe	ERR11029672	1.58e+06	13.66
RNA Illumina HiSeq 4000	ERR11040178	4.33e+07	6.54
RNA Illumina NovaSeq 6000	ERR12245545	9.15e+07	13.82

Nucleic acid extraction

The workflow for high molecular weight (HMW) DNA extraction at the Wellcome Sanger Institute (WSI) Tree of Life Core Laboratory includes a sequence of core procedures: sample preparation and homogenisation, DNA extraction, fragmentation and purification. Detailed protocols are available on protocols.io (Denton *et al.*, 2023). The fEleAnt2 sample was weighed and dissected on dry ice (Jay *et al.*, 2023), and tissue was cryogenically disrupted using the Covaris cryoPREP® Automated Dry Pulverizer (Narváez-Gómez *et al.*, 2023).

HMW DNA was extracted using the Automated MagAttract v1 protocol (Sheerin *et al.*, 2023). DNA was sheared into an average fragment size of 12–20 kb in a Megaruptor 3 system (Todorovic *et al.*, 2023). Sheared DNA was purified by solid-phase reversible immobilisation, using AMPure PB beads to eliminate shorter fragments and concentrate the DNA (Strickland *et al.*, 2023). The concentration of the sheared and purified DNA was assessed using a Nanodrop spectrophotometer and Qubit Fluorometer using the Qubit dsDNA High Sensitivity Assay kit. Fragment size distribution was evaluated by running the sample on the FemtoPulse system.

RNA was extracted from tissue of fEleAnt3 in the Tree of Life Laboratory at the WSI using the RNA Extraction: Automated MagMax™ mirVana protocol (do Amaral *et al.*, 2023). The RNA concentration was assessed using a Nanodrop spectrophotometer and a Qubit Fluorometer using the Qubit RNA Broad-Range Assay kit. Analysis of the integrity of the RNA was done using the Agilent RNA 6000 Pico Kit and Eukaryotic Total RNA assay.

Hi-C sample preparation

Hi-C data were generated from the tissue from the fEleAnt3 sample, using the Arima-HiC v2 kit. In brief, frozen tissue (stored at –80 °C) was fixed, and the DNA crosslinked using a TC buffer with 22% formaldehyde concentration. After crosslinking the tissue was homogenised using the Diagnocine Power Masher-II and BioMasher-II tubes and pestles. Following the kit manufacturer's instructions, crosslinked DNA was digested using a restriction enzyme master mix. The 5'-overhangs were then filled in and labelled with biotinylated nucleotides and proximally ligated. An overnight incubation was carried out for enzymes to digest remaining proteins and for crosslinks to reverse. A clean up was performed with SPRIselect beads prior to library preparation.

Table 2. Genome assembly data for *Electrona antarctica*, fEleAnt2.1.

Genome assembly		
Assembly name	fEleAnt2.1	
Assembly accession	GCA_951216825.1	
Accession of alternate haplotype	GCA_951214905.1	
Span (Mb)	1,427.40	
Number of contigs	5,080	
Number of scaffolds	1,900	
Longest scaffold (Mb)	65.73	
Assembly metrics*		Benchmark
Contig N50 length (Mb)	1.0	$\geq 1\text{ Mb}$
Scaffold N50 length (Mb)	53.6	= chromosome N50
Consensus quality (QV)	53.3	≥ 40
k-mer completeness	Primary: 74.95%; alternate: 71.26%; combined: 96.62%	$\geq 95\%$
BUSCO**	C:95.9%[S:83.1%,D:12.7%], F:1.1%,M:3.1%,n:3,640	$S > 90\%$, $D < 5\%$
Percentage of assembly mapped to chromosomes	88.72%	$\geq 90\%$
Sex chromosomes	Not identified	localised homologous pairs
Organelles	Mitochondrial genome: 20.02 kb	complete single alleles

* Assembly metric benchmarks are adapted from [Rhie et al. \(2021\)](#) and the Earth BioGenome Project Report on Assembly Standards [September 2024](#).

** BUSCO scores based on the actinopterygii_odb10 BUSCO set using version 5.3.2. C = complete [S = single copy, D = duplicated], F = fragmented, M = missing, n = number of orthologues in comparison. A full set of BUSCO scores is available at https://blobtoolkit.genomehubs.org/view/fEleAnt2_1/dataset/fEleAnt2_1/busco.

Library preparation and sequencing

PacBio HiFi

Libraries were prepared using the PacBio Express Template Preparation Kit v2.0 (Pacific Biosciences, California, USA) as per the manufacturer's instructions. The kit includes the reagents required for removal of single-strand overhangs, DNA damage repair, end repair/A-tailing, adapter ligation, and nuclease treatment. Library preparation also included a library purification step using AMPure PB beads (Pacific Biosciences, California, USA) and size selection step to remove templates shorter than 3 kb using AMPure PB modified SPRI. DNA concentration was quantified using the Qubit Fluorometer v2.0 (Thermo Fisher Scientific) and Qubit HS Assay Kit and the final library fragment size analysis was carried out using the Agilent Femto Pulse Automated Pulsed Field CE Instrument (Agilent Technologies).

Samples were sequenced using the Sequel IIe system (Pacific Biosciences, California, USA). The concentration of the library

loaded onto the Sequel IIe was in the range 40–135 pM. The SMRT link software, a PacBio web-based end-to-end workflow manager, was used to set-up and monitor the run, as well as perform primary and secondary analysis of the data upon completion.

Hi-C

For Hi-C library preparation, DNA was fragmented to a size of 400 to 600 bp using a Covaris E220 sonicator. The DNA was then enriched, barcoded, and amplified using the NEBNext Ultra II DNA Library Prep Kit following manufacturers' instructions. The Hi-C sequencing was performed using paired-end sequencing with a read length of 150 bp on an Illumina NovaSeq 6000.

RNA

RNA-Seq libraries were constructed using the NEB Ultra II RNA Library Prep kit, following the manufacturer's instructions. RNA sequencing was performed on the Illumina NovaSeq 6000 instrument.




Figure 2. Genome assembly of *Electrona antarctica*, fEleAnt2.1: metrics. The BlobToolKit snail plot shows N50 metrics and BUSCO gene completeness. The main plot is divided into 1,000 size-ordered bins around the circumference with each bin representing 0.1% of the 1,427,445,741 bp assembly. The distribution of scaffold lengths is shown in dark grey with the plot radius scaled to the longest scaffold present in the assembly (65,725,131 bp, shown in red). Orange and pale-orange arcs show the N50 and N90 scaffold lengths (53,633,514 and 1,105,000 bp), respectively. The pale grey spiral shows the cumulative scaffold count on a log scale with white scale lines showing successive orders of magnitude. The blue and pale-blue area around the outside of the plot shows the distribution of GC, AT and N percentages in the same bins as the inner plot. A summary of complete, fragmented, duplicated and missing BUSCO genes in the actinopterygii_odb10 set is shown in the top right. An interactive version of this figure is available at https://blobtoolkit.genomehubs.org/view/fEleAnt2_1/dataset/fEleAnt2_1/snail.

Genome assembly, curation and evaluation

Assembly

HiFi reads were assembled using Hifiasm (Cheng *et al.*, 2021) using the --primary option. Haplotypic duplications were identified and removed with purge_dups (Guan *et al.*, 2020). Hi-C reads were further mapped with bwa-mem2 (Vasimuddin *et al.*, 2019) to the primary contigs, which were further scaffolded using the provided Hi-C data (Rao *et al.*, 2014) in YaHS (Zhou *et al.*, 2023) using the --break option. Scaffolded assemblies were evaluated using Gfastats (Formenti *et al.*, 2022), BUSCO (Manni *et al.*, 2021) and MERQURY.FK (Rhie *et al.*, 2020).

The mitochondrial genome was assembled using MitoHiFi (Uliano-Silva *et al.*, 2023), which runs MitoFinder (Allio *et al.*, 2020) and uses these annotations to select the final mitochondrial contig and to ensure the general quality of the sequence.

Assembly curation

The assembly was decontaminated using the Assembly Screen for Cobionts and Contaminants (ASCC) pipeline (article in preparation). Manual curation was primarily conducted using PretextView (Harry, 2022), with additional insights provided by JBrowse2 (Diesh *et al.*, 2023) and HiGlass (Kerpedjiev *et al.*, 2018). Scaffolds were visually inspected and corrected as




Figure 3. Genome assembly of *Electrona antarctica*, fEleAnt2.1: BlobToolKit GC-coverage plot. Sequences are coloured by phylum. Circles are sized in proportion to sequence length. Histograms show the distribution of sequence length sum along each axis. An interactive version of this figure is available at https://blobtoolkit.genomehubs.org/view/fEleAnt2_1/dataset/fEleAnt2_1/blob.

described by [Howe *et al.* \(2021\)](#). Any identified contamination, missed joins, and mis-joins were corrected, and duplicate sequences were tagged and removed. The entire process is documented at <https://gitlab.com/wtsi-grit/rapid-curation> (article in preparation).

Evaluation of the final assembly

The Merqury.FK tool ([Rhie *et al.*, 2020](#)) was used to evaluate k -mer completeness and assembly quality for the primary and alternate haplotypes using the k -mer databases ($k = 31$) that were pre-computed prior to genome assembly. The analysis

outputs included assembly QV scores and completeness statistics.

A Hi-C contact map was produced for the final, public version of the assembly. The Hi-C reads were aligned using bwa-mem2 ([Vasimuddin *et al.*, 2019](#)) and the alignment files were combined using SAMtools ([Danecek *et al.*, 2021](#)). The Hi-C alignments were converted into a contact map using BEDTools ([Quinlan & Hall, 2010](#)) and the Cooler tool suite ([Abdennur & Mirny, 2020](#)). The contact map was visualised in HiGlass ([Kerpedjiev *et al.*, 2018](#)).




Figure 4. Genome assembly of *Electrona antarctica* fEleAnt2.1: BlobToolKit cumulative sequence plot. The grey line shows cumulative length for all sequences. Coloured lines show cumulative lengths of sequences assigned to each phylum using the buscogenes taxrule. An interactive version of this figure is available at https://blobtoolkit.genomehubs.org/view/fEleAnt2_1/dataset/fEleAnt2_1/cumulative.

The genome was analysed within the BlobToolKit environment (Challis *et al.*, 2020) and BUSCO completeness scores (Manni *et al.*, 2021) were calculated.

Table 4 contains a list of relevant software tool versions and sources.

Wellcome Sanger Institute – Legal and Governance

The materials that have contributed to this genome note have been supplied by a Tree of Life collaborator.

The Wellcome Sanger Institute employs a process whereby due diligence is carried out proportionate to the nature of the materials themselves, and the circumstances under which they have been/are to be collected and provided for use. The purpose of this is to address and mitigate any potential legal

and/or ethical implications of receipt and use of the materials as part of the research project, and to ensure that in doing so we align with best practice wherever possible.

The overarching areas of consideration are:

- Ethical review of provenance and sourcing of the material
- Legality of collection, transfer and use (national and international)

Each transfer of samples is undertaken according to a Research Collaboration Agreement or Material Transfer Agreement entered into by the Tree of Life collaborator, Genome Research Limited (operating as the Wellcome Sanger Institute) and in some circumstances other Tree of Life collaborators.




Figure 5. Genome assembly of *Electrona antarctica* fEleAnt2.1: Hi-C contact map of the fEleAnt2.1 assembly, visualised using HiGlass. Chromosomes are shown in order of size from left to right and top to bottom. An interactive version of this figure may be viewed at <https://genome-note-higlass.tol.sanger.ac.uk/I/?d=Bmyjwb5DRka--g6D5GDvUA>.

Table 3. Chromosomal pseudomolecules in the genome assembly of *Electrona antarctica*, fEleAnt2.

INSDC accession	Name	Length (Mb)	GC%
OX578246.1	1	65.73	43.5
OX578247.1	2	64.11	43.0
OX578248.1	3	63.62	43.5
OX578249.1	4	62.16	44.0
OX578250.1	5	60.55	43.5
OX578251.1	6	60.31	43.5
OX578252.1	7	59.04	43.5
OX578253.1	8	58.87	43.0
OX578254.1	9	58.47	43.5
OX578255.1	10	56.06	43.5
OX578256.1	11	54.82	43.0

INSDC accession	Name	Length (Mb)	GC%
OX578257.1	12	53.63	43.0
OX578258.1	13	52.08	43.5
OX578259.1	14	50.67	43.5
OX578260.1	15	49.8	43.5
OX578261.1	16	48.97	44.0
OX578262.1	17	48.42	43.5
OX578263.1	18	48.16	43.5
OX578264.1	19	44.95	43.0
OX578265.1	20	43.09	43.5
OX578266.1	21	42.9	43.5
OX578267.1	22	41.92	43.5
OX578268.1	23	39.46	43.5
OX578269.1	24	34.7	44.0
OX578270.1	MT	0.02	48.5

Table 4. Software tools: versions and sources.

Software tool	Version	Source
BEDTools	2.30.0	https://github.com/arq5x/bedtools2
BLAST	2.14.0	ftp://ftp.ncbi.nlm.nih.gov/blast/executables/blast+/
BlobToolKit	4.3.7	https://github.com/blobtoolkit/blobtoolkit
BUSCO	5.4.3 and 5.5.0	https://gitlab.com/ezlab/busco
bwa-mem2	2.2.1	https://github.com/bwa-mem2/bwa-mem2
Cooler	0.8.11	https://github.com/open2c/cooler
FastK	427104ea91c78c3b8b8b49f1a7d6bbeaa869ba1c	https://github.com/thegenemyers/FASTK
Gfastats	1.3.6	https://github.com/vgl-hub/gfastats
Hifiasm	0.19.8-r587	https://github.com/chhylp123/hifiasm
HiGlass	44086069ee7d4d3f6f3f0012569789ec138f42b84a a44357826c0b6753eb28de	https://github.com/higlass/higlass
Mercury.FK	d00d98157618f4e8d1a9190026b19b471055b22e	https://github.com/thegenemyers/MERQURY.FK
MitoHiFi	3	https://github.com/marcelauliano/MitoHiFi
Nextflow	23.04.0-5857	https://github.com/nextflow-io/nextflow
PretextView	0.2.5	https://github.com/sanger-tol/PretextView
purge_dups	1.2.5	https://github.com/dfguan/purge_dups
samtools	1.16.1, 1.17, and 1.18	https://github.com/samtools/samtools
sanger-tol/ascc	-	https://github.com/sanger-tol/ascc
Singularity	3.9.0	https://github.com/sylabs/singularity
YaHS	1.2a.2	https://github.com/c-zhou/yahs

Data availability

European Nucleotide Archive: *Electrona antarctica* (Antarctic lanternfish). Accession number PRJEB60649; <https://identifiers.org/ena.embl/PRJEB60649>. The genome sequence is released openly for reuse. The *Electrona antarctica* genome sequencing initiative is part of the Vertebrate Genomes Project (PRJNA489243). All raw sequence data and the assembly have been deposited in INSDC databases. The genome will be annotated using available RNA-Seq data and presented through the [Ensembl](#) pipeline at the European Bioinformatics Institute. Raw data and assembly accession identifiers are reported in [Table 1](#) and [Table 2](#).

Author information

Members of the Wellcome Sanger Institute Tree of Life Management, Samples and Laboratory team are listed here: <https://doi.org/10.5281/zenodo.12162482>.

Members of Wellcome Sanger Institute Scientific Operations: Sequencing Operations are listed here: <https://doi.org/10.5281/zenodo.12165051>.

Members of the Wellcome Sanger Institute Tree of Life Core Informatics team are listed here: <https://doi.org/10.5281/zenodo.12160324>.

Members of the Tree of Life Core Informatics collective are listed here: <https://doi.org/10.5281/zenodo.12205391>.

Acknowledgements

We thank the crew and science team of the James Clark Ross research vessel and the British Antarctic Survey logistics team for their support during the JR19001 research cruise.

References

- Abdennur N, Mirny LA: **Cooler: scalable storage for Hi-C data and other genomically labeled arrays.** *Bioinformatics.* 2020; **36**(1): 311–316.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Allio R, Schomaker-Bastos A, Romiguier J, *et al.*: **MitoFinder: efficient automated large-scale extraction of mitogenomic data in target enrichment phylogenomics.** *Mol Ecol Resour.* 2020; **20**(4): 892–905.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Catul V, Gauns M, Karuppasamy PK: **A review on mesopelagic fishes belonging to family Myctophidae.** *Rev Fish Biol Fisheries.* 2011; **21**(3): 339–354.
[Publisher Full Text](#)
- Challis R, Richards E, Rajan J, *et al.*: **BlobToolKit – interactive quality assessment of genome assemblies.** *G3 (Bethesda).* 2020; **10**(4): 1361–1374.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Cheng H, Concepcion GT, Feng X, *et al.*: **Haplotype-resolved *de novo* assembly using phased assembly graphs with hifiasm.** *Nat Methods.* 2021; **18**(2): 170–175.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Collins MA, Stowasser G, Fielding S, *et al.*: **Latitudinal and bathymetric patterns in the distribution and abundance of mesopelagic fish in the Scotia Sea.** *Deep-Sea Res II: Top Stud Oceanogr.* 2012; **59–60**: 189–198.
[Publisher Full Text](#)
- Danecek P, Bonfield JK, Liddle J, *et al.*: **Twelve years of SAMtools and BCFtools.** *GigaScience.* 2021; **10**(2): giab008.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Denton A, Yatsenko H, Jay J, *et al.*: **Sanger Tree of Life wet laboratory protocol collection V.1** *protocols.io.* 2023.
[Publisher Full Text](#)
- Diesh C, Stevens GJ, Xie P, *et al.*: **JBrowse 2: a modular genome browser with views of synteny and structural variation.** *Genome Biol.* 2023; **24**(1): 74.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- do Amaral RJV, Denton A, Yatsenko H, *et al.*: **Sanger Tree of Life RNA extraction: automated MagMax™ mirVana.** *protocols.io.* 2023.
[Publisher Full Text](#)
- Formenti G, Abueg L, Brajuka A, *et al.*: **Gfstats: conversion, evaluation and manipulation of genome sequences using assembly graphs.** *Bioinformatics.* 2022; **38**(17): 4214–4216.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Guan D, McCarthy SA, Wood J, *et al.*: **Identifying and removing haplotypic duplication in primary genome assemblies.** *Bioinformatics.* 2020; **36**(9): 2896–2898.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Harry E: **PretextView (Paired REad TEXTure Viewer): a desktop application for viewing pretext contact maps.** 2022.
[Reference Source](#)
- Howe K, Chow W, Collins J, *et al.*: **Significantly improving the quality of genome assemblies through curation.** *GigaScience.* 2021; **10**(1): g1aa153.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Hulley PA: **Myctophidae.** In: Gon O. and Heemstra, P. C. (eds.) *Fishes of the Southern Ocean.* Grahamstown, South Africa: J.L.B. Smith Institute of Ichthyology, 1990; 146–178.
- Hulley PA, Duhamel G: **Aspects of lanternfish distribution in the Kerguelen Plateau region.** In: Duhamel, G. and Welsford, D. (eds.) *The Kerguelen Plateau: Marine Ecosystems and Fisheries.* Paris: Société Française d'Ichtyologie, 2011; **35**(SP): 183–195.
[Publisher Full Text](#)
- Jay J, Yatsenko H, Narváez-Gómez JP, *et al.*: **Sanger Tree of Life sample preparation: triage and dissection.** *protocols.io.* 2023.
[Publisher Full Text](#)
- Kerpedjiev P, Abdennur N, Lekschas F, *et al.*: **HiGlass: web-based visual exploration and analysis of genome interaction maps.** *Genome Biol.* 2018; **19**(1): 125.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Manni M, Berkeley MR, Seppey M, *et al.*: **BUSCO update: novel and streamlined workflows along with broader and deeper phylogenetic coverage for scoring of eukaryotic, prokaryotic, and viral genomes.** *Mol Biol Evol.* 2021; **38**(10): 4647–4654.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Narváez-Gómez JP, Mbye H, Oatley G, *et al.*: **Sanger Tree of Life sample homogenisation: Covaris cryoPREP® Automated Dry Pulverizer V.1.** *protocols.io.* 2023.
[Publisher Full Text](#)
- Quinlan AR, Hall IM: **BEDTools: a flexible suite of utilities for comparing genomic features.** *Bioinformatics.* 2010; **26**(6): 841–842.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Rao SSP, Huntley MH, Durand NC, *et al.*: **A 3D map of the human genome at kilobase resolution reveals principles of chromatin looping.** *Cell.* 2014; **159**(7): 1665–1680.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Rhie A, McCarthy SA, Fedrigo O, *et al.*: **Towards complete and error-free genome assemblies of all vertebrate species.** *Nature.* 2021; **592**(7856): 737–746.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Rhie A, Walenz BP, Koren S, *et al.*: **Merquy: reference-free quality, completeness, and phasing assessment for genome assemblies.** *Genome Biol.* 2020; **21**(1): 245.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Sheerin E, Sampaio F, Oatley G: **Sanger Tree of Life HMW DNA extraction: automated MagAttract v.1.** *protocols.io.* 2023.
[Publisher Full Text](#)
- Strickland M, Cornwell C, Howard C: **Sanger Tree of Life fragmented DNA clean up: manual SPRI.** *protocols.io.* 2023.
[Publisher Full Text](#)
- Todorovic M, Sampaio F, Howard C: **Sanger Tree of Life HMW DNA fragmentation: diagenode Megaruptor³ for PacBio HiFi.** *protocols.io.* 2023.
[Publisher Full Text](#)
- Uliano-Silva M, Ferreira JGRN, Krashenninnikova K, *et al.*: **MitoHiFi: a python pipeline for mitochondrial genome assembly from PacBio high fidelity reads.** *BMC Bioinformatics.* 2023; **24**(1): 288.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Vasimuddin M, Misra S, Li H, *et al.*: **Efficient architecture-aware acceleration of BWA-MEM for multicore systems.** In: *2019 IEEE International Parallel and Distributed Processing Symposium (IPDPS).* IEEE, 2019; 314–324.
[Publisher Full Text](#)
- Williams A: **Electrona antarctica.** 2019.
[Publisher Full Text](#)
- Zhou C, McCarthy SA, Durbin R: **YaHS: Yet another Hi-C Scaffolding tool.** *Bioinformatics.* 2023; **39**(1): btac808.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Open Peer Review

Current Peer Review Status:   

Version 2

Reviewer Report 12 August 2025

<https://doi.org/10.21956/wellcomeopenres.27236.r127706>

© 2025 Whibley A. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Annabel Whibley 

Grapevine Improvement, Bragato Research Institute, Lincoln, New Zealand

The authors have corrected the reporting error. Thank you. I have no further comments.

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Genomics Bioinformatics

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Version 1

Reviewer Report 17 March 2025

<https://doi.org/10.21956/wellcomeopenres.26253.r119700>

© 2025 Whibley A. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Annabel Whibley 

Grapevine Improvement, Bragato Research Institute, Lincoln, New Zealand

Bista, Collins and colleagues report the genome assembly of the Antarctic lanternfish, *Electrona antarctica*. Relative to other DTOL Genome Notes that I have reviewed, this assembly is more highly fragmented and with slightly inferior quality statistics, presumably reflecting challenges from the sample properties. It is nevertheless a high- quality assembly, with the primary assembly

showing good completeness and accuracy and being robustly scaffolded into 24 pseudomolecules.

I believe there is an error in the reporting of the total of long read sequence data that went into the initial assembly (13.66Gb data is not 43x fold coverage of a 1.4Gb genome). Reviewing the information in Table1, I think the authors have reported coverage for just one of the Sequel IIe runs, not all of them.

The data note is well-written, with clear reporting of metadata and methods, following the DTOL template. The data accessions are functional. RNA sequence data was obtained but an annotation has not yet been reported.

Is the rationale for creating the dataset(s) clearly described?

Yes

Are the protocols appropriate and is the work technically sound?

Yes

Are sufficient details of methods and materials provided to allow replication by others?

Yes

Are the datasets clearly presented in a useable and accessible format?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Genomics Bioinformatics

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Reviewer Report 10 March 2025

<https://doi.org/10.21956/wellcomeopenres.26253.r119695>

© 2025 Senevirathna J. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Jayan Duminda M. Senevirathna 

Uva Wellassa University, Badulla, Uva Province, Sri Lanka

The study is perfect and well written in this article. The data are sufficient and nicely presented with appropriate results and publicly available.

Hence, I would like to accept the manuscript for publication with minor modifications. Followings are suggested for verifications or improvements;

- In Hi-C sample preparation, the frozen tissue was fixed, with what...?
- Table 2, and figure 3 titles are quiet similar.. better to change...
- "The total count of gaps in the scaffolds is 3,179." this sentence is not clear.. or maybe not necessary
- Why you could not identify sex chromosomes, XX..?
- In keywords; *Electrona antarctica* should be italic.. and keep consistency

Is the rationale for creating the dataset(s) clearly described?

Yes

Are the protocols appropriate and is the work technically sound?

Yes

Are sufficient details of methods and materials provided to allow replication by others?

Yes

Are the datasets clearly presented in a useable and accessible format?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Aquatic Molecular Biology and Biotechnology

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 06 March 2025

<https://doi.org/10.21956/wellcomeopenres.26253.r119699>

© 2025 Kishida T. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Takushi Kishida 

Nihon University, Fujisawa, Japan

The authors provide a chromosomal-level phased genome assembly of an Antarctic lanternfish. This genome assembly is of high quality, and I have no more suggestions about their methodologies.

I have only one comment about their specimens. I strongly recommend that the authors deposit the specimens used in this study (fEleAnt2 and fEleAnt3) in an appropriate museum. I think high-quality genomic data should be accompanied by vouchered specimens.

Is the rationale for creating the dataset(s) clearly described?

Yes

Are the protocols appropriate and is the work technically sound?

Yes

Are sufficient details of methods and materials provided to allow replication by others?

Partly

Are the datasets clearly presented in a useable and accessible format?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Genomics, Population genetics

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.
