



DATA NOTE

The genome sequence of the Brindled White-spot moth,

Parectropis similaria (Hufnagel, 1767)

[version 1; peer review: 4 approved]

Douglas Boyes¹⁺, Gavin R. Broad^{id}², David C. Lees^{id}², Clare Boyes³,
Natural History Museum Genome Acquisition Lab,
Darwin Tree of Life Barcoding collective,
Wellcome Sanger Institute Tree of Life Management, Samples and Laboratory
team,
Wellcome Sanger Institute Scientific Operations: Sequencing Operations,
Wellcome Sanger Institute Tree of Life Core Informatics team,
Tree of Life Core Informatics collective, Darwin Tree of Life Consortium

¹UK Centre for Ecology & Hydrology, Wallingford, England, UK

²Natural History Museum, London, England, UK

³Independent researcher, Welshpool, Wales, UK

+ Deceased author

V1 First published: 08 Apr 2025, 10:179
<https://doi.org/10.12688/wellcomeopenres.23984.1>
Latest published: 08 Apr 2025, 10:179
<https://doi.org/10.12688/wellcomeopenres.23984.1>

Abstract

We present a genome assembly from a female *Parectropis similaria* (Brindled White-spot; Arthropoda; Insecta; Lepidoptera; Geometridae). The genome sequence has a total length of 574.79 megabases. Most of the assembly (99.92%) is scaffolded into 31 chromosomal pseudomolecules, including the Z sex chromosome. The mitochondrial genome has also been assembled, with a length of 16.94 kilobases.

Keywords

Parectropis similaria, Brindled White-spot, genome sequence, chromosomal, Lepidoptera



This article is included in the [Tree of Life gateway](#).

Open Peer Review

Approval Status

	1	2	3	4
version 1				
08 Apr 2025	view	view	view	view

1. **Laurence Despres** ^{id}, Laboratoire d'Ecologie Alpine (Ringgold ID: 56837), Grenoble, France
2. **Annabel Whibley** ^{id}, The University of Auckland, Auckland, New Zealand
3. **Panagiotis Ioannidis** ^{id}, Foundation for Research & Technology - Hellas, Crete, Greece
4. **Arjen Van 't Hof** ^{id}, Biology Centre of the Czech Academy of Sciences, Budějovice, Czech Republic

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding author: Darwin Tree of Life Consortium (mark.blaxter@sanger.ac.uk)

Author roles: **Boyes D:** Investigation, Resources; **Broad GR:** Investigation, Resources; **Lees DC:** Investigation, Resources; **Boyes C:** Writing – Original Draft Preparation;

Competing interests: No competing interests were disclosed.

Grant information: This work was supported by Wellcome through core funding to the Wellcome Sanger Institute (220540) and the Darwin Tree of Life Discretionary Award [218328, <https://doi.org/10.35802/218328>].

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2025 Boyes D *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Boyes D, Broad GR, Lees DC *et al.* **The genome sequence of the Brindled White-spot moth, *Parectropis similaria* (Hufnagel, 1767) [version 1; peer review: 4 approved]** Wellcome Open Research 2025, 10:179 <https://doi.org/10.12688/wellcomeopenres.23984.1>

First published: 08 Apr 2025, 10:179 <https://doi.org/10.12688/wellcomeopenres.23984.1>

Species taxonomy

Eukaryota; Opisthokonta; Metazoa; Eumetazoa; Bilateria; Protostomia; Ecdysozoa; Panarthropoda; Arthropoda; Mandibulata; Pancrustacea; Hexapoda; Insecta; Dicondylia; Pterygota; Neoptera; Endopterygota; Amphiesmenoptera; Lepidoptera; Glossata; Neolepidoptera; Heteroneura; Ditrysia; Obtectomera; Geometroidea; Geometridae; Ennominae; *Parectropis*; *Parectropis similaria* (Hufnagel, 1767) (NCBI:txid934885)

Background

The Brindled White-spot (*Parectropis similaria*) is a moth in the family Geometridae. It is a moth of ancient woodlands, and is locally common in southern and eastern England. It has declined significantly since 1970 (Randle *et al.*, 2019). It is a meso-thermophilous species distributed through the Palaearctic region, from Britain in the west to Japan in the east (GBIF Secretariat, 2023).

As its common name suggests, the moth's appearance is brindled and, particularly in fresh specimens, sports a large pale patch near the outer edge of the forewing. There is one generation a year, flying from late May into June. The caterpillars feed on a range of broad-leaved trees, but particularly birch, hawthorn and oak (Waring *et al.*, 2017). The caterpillar has two colour forms – a brown and a green. It pupates underground where it spends the winter (Henwood *et al.*, 2020).

We present a chromosome-level genome sequence for *Parectropis similaria* based on a female specimen from Wytham Woods, Oxfordshire, UK.

Genome sequence report

Sequencing data

The genome of a specimen of *Parectropis similaria* (Figure 1) was sequenced using Pacific Biosciences single-molecule HiFi long reads, generating 24.93 Gb (gigabases) from 2.02 million reads. GenomeScope analysis of the PacBio HiFi data estimated the haploid genome size at 557.31 Mb, with a heterozygosity of 0.83% and repeat content of 30.40%. These values provide an initial assessment of genome complexity



Figure 1. Photograph of the *Parectropis similaria* (ilParSimi1) specimen used for genome sequencing.

and the challenges anticipated during assembly. Based on this estimated genome size, the sequencing data provided approximately 42.0x coverage of the genome. Chromosome conformation Hi-C sequencing produced 108.32 Gb from 717.35 million reads. Table 1 summarises the specimen and sequencing information.

Assembly statistics

The primary haplotype was assembled, and contigs corresponding to an alternate haplotype were also deposited in INSDC databases. The assembly was improved by manual curation, which corrected 19 misjoins or missing joins and removed four haplotypic duplications. The final assembly has a total length of 574.79 Mb in 36 scaffolds, with 69 gaps, and a scaffold N50 of 20.85 Mb (Table 2).

The snail plot in Figure 2 provides a summary of the assembly statistics, indicating the distribution of scaffold lengths and other assembly metrics. Figure 3 shows the distribution of scaffolds by GC proportion and coverage. Figure 4 presents a cumulative assembly plot, with separate curves representing different scaffold subsets assigned to various phyla, illustrating the completeness of the assembly.

Most of the assembly sequence (99.94%) was assigned to 31 chromosomal-level scaffolds, representing 30 autosomes and the Z sex chromosome. These chromosome-level scaffolds, confirmed by Hi-C data, are named according to size (Figure 5; Table 3). During curation, the Z chromosome was assigned based on synteny to the genome of *Agriopis aurantiaria* (GCA_914767915.1) (Boyce *et al.*, 2023). This species appears to exhibit the ZO karyotype.

The mitochondrial genome was also assembled. This sequence is included as a contig in the multifasta file of the genome submission and as a standalone record.

Assembly quality metrics

The estimated Quality Value (QV) and *k*-mer completeness metrics, along with BUSCO completeness scores, were calculated for each haplotype and the combined assembly. The QV reflects the base-level accuracy of the assembly, while *k*-mer completeness indicates the proportion of expected *k*-mers identified in the assembly. BUSCO scores provide a measure of completeness based on benchmarking universal single-copy orthologues.

The combined primary and alternate assemblies achieve an estimated QV of 65.8. The *k*-mer recovery for the primary haplotype is 83.53%, for the alternate haplotype 76.07%, and the combined primary and alternate assemblies have a *k*-mer recovery of 99.23%. BUSCO analysis using the lepidoptera_odb10 reference set (*n* = 5,286) identified 98.5% of the expected gene set (single = 98.1%, duplicated = 0.5%).

Table 2 provides assembly metric benchmarks adapted from Rhie *et al.* (2021) and the Earth BioGenome Project (EBP) Report on Assembly Standards September 2024. The assembly achieves the EBP reference standard of 7.C.Q65.

Table 1. Specimen and sequencing data for *Parectropis similaria*.

Project information			
Study title	Parectropis similaria (brindled white spot)		
Umbrella BioProject	PRJEB71300		
Species	<i>Parectropis similaria</i>		
BioSpecimen	SAMEA10979183		
NCBI taxonomy ID	934885		
Specimen information			
Technology	ToLID	BioSample accession	Organism part
PacBio long read sequencing	ilParSimi1	SAMEA10979620	head and thorax
Hi-C sequencing	ilParSimi2	SAMEA114805714	whole organism
RNA sequencing	ilParSimi2	SAMEA114805714	whole organism
Sequencing information			
Platform	Run accession	Read count	Base count (Gb)
Hi-C Illumina NovaSeq X	ERR13702754	7.17e+08	108.32
PacBio Sequel IIe	ERR12370426	2.02e+06	24.93
RNA Illumina NovaSeq X	ERR13999092	9.77e+07	14.76

Methods

Sample acquisition and DNA barcoding

An adult female *Parectropis similaria* (specimen ID Ox001920, ToLID ilParSimi1) was collected from Wytham Woods, Oxfordshire, United Kingdom (latitude 51.77, longitude -1.34) on 2021-06-16 by light trap. The specimen was collected and identified by Douglas Boyes (University of Oxford) and preserved on dry ice.x

The specimen used for Hi-C and RNA sequencing (specimen ID NHMUK010881093, ToLID ilParSimi2) was collected from Tudeley Woods, England, United Kingdom (latitude 51.16, longitude 0.31) on 2022-06-10. The specimen was collected by Gavin Broad (Natural History Museum) and identified by David Lees (Natural History Museum) and preserved by dry freezing (-80 °C).

The initial identification by Expert Id was verified by an additional DNA barcoding process according to the framework developed by Twyford *et al.* (2024). A small sample was dissected from the specimen and stored in ethanol, while the remaining parts were shipped on dry ice to the Wellcome Sanger Institute (WSI) (Pereira *et al.*, 2022). The tissue was lysed, the COI marker region was amplified by PCR, and amplicons were sequenced and compared to the BOLD database, confirming the species identification (Crowley *et al.*, 2023). Following whole genome sequence generation, the relevant DNA barcode region was also used alongside the initial barcoding data for

sample tracking at the WSI (Twyford *et al.*, 2024). The standard operating procedures for Darwin Tree of Life barcoding have been deposited on protocols.io (Beasley *et al.*, 2023).

Metadata collection for samples adhered to the Darwin Tree of Life project standards described by Lawniczak *et al.* (2022).

Nucleic acid extraction

The workflow for high molecular weight (HMW) DNA extraction at the Wellcome Sanger Institute (WSI) Tree of Life Core Laboratory includes a sequence of procedures: sample preparation and homogenisation, DNA extraction, fragmentation and purification. Detailed protocols are available on protocols.io (Denton *et al.*, 2023b). The ilParSimi1 sample was prepared for DNA extraction by weighing and dissecting it on dry ice (Jay *et al.*, 2023). Tissue from the head and thorax was homogenised using a PowerMasher II tissue disruptor (Denton *et al.*, 2023a).

HMW DNA was extracted in the WSI Scientific Operations core using the Automated MagAttract v2 protocol (Oatley *et al.*, 2023). The DNA was sheared into an average fragment size of 12–20 kb in a Megaruptor 3 system (Bates *et al.*, 2023). Sheared DNA was purified by solid-phase reversible immobilisation, using AMPure PB beads to eliminate shorter fragments and concentrate the DNA (Strickland *et al.*, 2023). The concentration of the sheared and purified DNA was assessed using a Nanodrop spectrophotometer and Qubit Fluorometer

Table 2. Genome assembly data for *Parectropis similaria*.

Genome assembly		
Assembly name	ilParSimi1.1	
Assembly accession	GCA_964276655.1	
Alternate haplotype accession	GCA_964276675.1	
Assembly level for primary assembly	chromosome	
Span (Mb)	574.79	
Number of contigs	105	
Number of scaffolds	36	
Longest scaffold (Mb)	26.48	
Assembly metric	Measure	Benchmark
Contig N50 length	10.63 Mb	≥ 1 Mb
Scaffold N50 length	20.85 Mb	= chromosome N50
Consensus quality (QV)	Primary: 66.0; alternate: 65.7; combined: 65.8	≥ 40
k-mer completeness	Primary: 83.53%; alternate: 76.07%; combined: 99.23%	$\geq 95\%$
BUSCO*	C:98.5%[S:98.1%,D:0.5%], F:0.2%,M:1.2%,n:5,286	$S > 90\%$; $D < 5\%$
Percentage of assembly mapped to chromosomes	99.94%	$\geq 90\%$
Sex chromosomes	Z	localised homologous pairs
Organelles	Mitochondrial genome: 16.94 kb	complete single alleles

* BUSCO scores based on the lepidoptera_odb10 BUSCO set using version 5.5.0. C = complete [S = single copy, D = duplicated], F = fragmented, M = missing, n = number of orthologues in comparison.

using the Qubit dsDNA High Sensitivity Assay kit. Fragment size distribution was evaluated by running the sample on the FemtoPulse system.

RNA was extracted from whole organism tissue of ilParSimi2 in the Tree of Life Laboratory at the WSI using the RNA Extraction: Automated MagMax™ mirVana protocol (do Amaral *et al.*, 2023). The RNA concentration was assessed using a Nanodrop spectrophotometer and a Qubit Fluorometer using the Qubit RNA Broad-Range Assay kit. Analysis of the integrity of the RNA was done using the Agilent RNA 6000 Pico Kit and Eukaryotic Total RNA assay.

Hi-C sample preparation

Tissue from the whole organism of the ilParSimi2 sample was processed for Hi-C sequencing at the WSI Scientific Operations core, using the Arima-HiC v2 kit. In brief, 20–50 mg of frozen tissue (stored at -80°C) was fixed, and the DNA crosslinked using a TC buffer with 22% formaldehyde concentration. After crosslinking, the tissue was homogenised using the

Diagnocine Power Masher-II and BioMasher-II tubes and pestles. Following the Arima-HiC v2 kit manufacturer's instructions, crosslinked DNA was digested using a restriction enzyme master mix. The 5'-overhangs were filled in and labelled with biotinylated nucleotides and proximally ligated. An overnight incubation was carried out for enzymes to digest remaining proteins and for crosslinks to reverse. A clean up was performed with SPRIselect beads prior to library preparation. Additionally, the biotinylation percentage was estimated using the Qubit Fluorometer v4.0 (Thermo Fisher Scientific) and Qubit HS Assay Kit and Arima-HiC v2 QC beads.

Library preparation and sequencing

Library preparation and sequencing were performed at the WSI Scientific Operations core.

PacBio HiFi

At a minimum, samples were required to have an average fragment size exceeding 8 kb and a total mass over 400 ng to proceed to the low input SMRTbell Prep Kit 3.0 protocol (Pacific




Figure 2. Genome assembly of *Parectropis similaria*, ilParSimi1.1: metrics. The BlobToolKit snail plot provides an overview of assembly metrics and BUSCO gene completeness. The circumference represents the length of the whole genome sequence, and the main plot is divided into 1,000 bins around the circumference. The outermost blue tracks display the distribution of GC, AT, and N percentages across the bins. Scaffolds are arranged clockwise from longest to shortest and are depicted in dark grey. The longest scaffold is indicated by the red arc, and the deeper orange and pale orange arcs represent the N50 and N90 lengths. A light grey spiral at the centre shows the cumulative scaffold count on a logarithmic scale. A summary of complete, fragmented, duplicated, and missing BUSCO genes in the lepidoptera_odb10 set is presented at the top right. An interactive version of this figure is available at https://blobtoolkit.genomehubs.org/view/GCA_964276655.1/snail.

Biosciences, California, USA), depending on genome size and sequencing depth required. Libraries were prepared using the SMRTbell Prep Kit 3.0 (Pacific Biosciences, California, USA) as per the manufacturer's instructions. The kit includes the reagents required for end repair/A-tailing, adapter ligation, post-ligation SMRTbell bead cleanup, and nuclease treatment. Following the manufacturer's instructions, size selection and clean up was carried out using diluted AMPure PB beads (Pacific Biosciences, California, USA). DNA concentration was quantified using the Qubit Fluorometer v4.0 (Thermo Fisher Scientific) with Qubit 1X dsDNA HS assay kit and the final library fragment size analysis was carried out using the Agilent Femto Pulse Automated Pulsed Field CE Instrument (Agilent Technologies) and gDNA 55kb BAC analysis kit.

Samples were sequenced using the Sequel IIe system (Pacific Biosciences, California, USA). The concentration of the library loaded onto the Sequel IIe was in the range 40–135 pM. The SMRT link software, a PacBio web-based end-to-end workflow manager, was used to set-up and monitor the run, as well as perform primary and secondary analysis of the data upon completion.

Hi-C

For Hi-C library preparation, DNA was fragmented using the Covaris E220 sonicator (Covaris) and size selected using SPRISelect beads to 400 to 600 bp. The DNA was then enriched using the Arima-HiC v2 kit Enrichment beads. Using the NEBNext Ultra II DNA Library Prep Kit (New England




Figure 3. Genome assembly of *Parectropis similaria*, ilParSimi1.1: BlobToolKit GC-coverage plot. Blob plot showing sequence coverage (vertical axis) and GC content (horizontal axis). The circles represent scaffolds, with the size proportional to scaffold length and the colour representing phylum membership. The histograms along the axes display the total length of sequences distributed across different levels of coverage and GC content. An interactive version of this figure is available at https://blobtoolkit.genomehubs.org/view/GCA_964276655.1/blob.

Biolabs) for end repair, a-tailing, and adapter ligation. This uses a custom protocol which resembles the standard NEBNext Ultra II DNA Library Prep protocol but where library preparation occurs while DNA is bound to the Enrichment beads. For library amplification, 10 to 16 PCR cycles were required, determined by the sample biotinylation percentage. The Hi-C sequencing was performed using paired-end sequencing with a read length of 150 bp on an Illumina NovaSeq X instrument.

RNA

Poly(A) RNA-Seq libraries were constructed using the NEB Ultra II RNA Library Prep kit, following the manufacturer's instructions. RNA sequencing was performed on the Illumina NovaSeq X instrument.

Genome assembly, curation and evaluation

Assembly

Prior to assembly of the PacBio HiFi reads, a database of k -mer counts ($k = 31$) was generated from the filtered reads

using FastK. GenomeScope2 (Ranallo-Benavidez *et al.*, 2020) was used to analyse the k -mer frequency distributions, providing estimates of genome size, heterozygosity, and repeat content.

The HiFi reads were first assembled using Hifiasm (Cheng *et al.*, 2021) with the --primary option. The Hi-C reads were mapped to the primary contigs using bwa-mem2 (Vasimuddin *et al.*, 2019). The contigs were further scaffolded using the provided Hi-C data (Rao *et al.*, 2014) in YaHS (Zhou *et al.*, 2023) using the --break option for handling potential misassemblies. The scaffolded assemblies were evaluated using Gfastats (Formenti *et al.*, 2022), BUSCO (Manni *et al.*, 2021) and MERQUY.FK (Rhie *et al.*, 2020).

The mitochondrial genome was assembled using MitoHiFi (Uliano-Silva *et al.*, 2023), which runs MitoFinder (Allio *et al.*, 2020) and uses these annotations to select the final mitochondrial contig and to ensure the general quality of the sequence.




Figure 4. Genome assembly of *Parectropis similaria*, ilParSimi1.1: BlobToolKit cumulative sequence plot. The grey line shows cumulative length for all scaffolds. Coloured lines show cumulative lengths of scaffolds assigned to each phylum using the buscogenes taxrule. An interactive version of this figure is available at https://blobtoolkit.genomehubs.org/view/GCA_964276655.1/cumulative.

Assembly curation

The assembly was decontaminated using the Assembly Screen for Cobionts and Contaminants (ASCC) pipeline. Flat files and maps used in curation were generated via the TreeVal pipeline (Pointon *et al.*, 2023). Manual curation was conducted primarily in PretextView (Harry, 2022) and HiGlass (Kerpedjiev *et al.*, 2018), with additional insights provided by JBrowse2 (Diesh *et al.*, 2023). Scaffolds were visually inspected and corrected as described by Howe *et al.* (2021). Any identified contamination, missed joins, and mis-joins were amended, and duplicate sequences were tagged and removed. The sex chromosome was identified by synteny analysis. The curation process is documented at <https://gitlab.com/wtsi-grit/rapid-curation>.

Assembly quality assessment

The Merqury.FK tool (Rhie *et al.*, 2020), run in a Singularity container (Kurtzer *et al.*, 2017), was used to evaluate *k*-mer completeness and assembly quality for the primary and alternate haplotypes using the *k*-mer databases (*k* = 31) that were

computed prior to genome assembly. The analysis outputs included assembly QV scores and completeness statistics.

The blobtoolkit pipeline is a Nextflow port of the previous Snakemake Blobtoolkit pipeline (Challis *et al.*, 2020). It aligns the PacBio reads in SAMtools (Danecek *et al.*, 2021) and minimap2 (Li, 2018) and generates coverage tracks for regions of fixed size. In parallel, it queries the GoAT database (Challis *et al.*, 2023) to identify all matching BUSCO lineages to run BUSCO (Manni *et al.*, 2021). For the three domain-level BUSCO lineages, the pipeline aligns the BUSCO genes to the UniProt Reference Proteomes database (Bateman *et al.*, 2023) with DIAMOND blastp (Buchfink *et al.*, 2021). The genome is also divided into chunks according to the density of the BUSCO genes from the closest taxonomic lineage, and each chunk is aligned to the UniProt Reference Proteomes database using DIAMOND blastx. Genome sequences without a hit are chunked using seqtk and aligned to the NT database with blastn (Altschul *et al.*, 1990). The blobtools suite combines all these outputs into a blobdir for visualisation.




Figure 5. Genome assembly of *Parectropis similaria*, ilParSimi1.1: Hi-C contact map of the ilParSimi1.1 assembly, produced in PretextView. Chromosomes are shown in order of size from left to right and top to bottom.

Table 3. Chromosomal pseudomolecules in the genome assembly of *Parectropis similaria*, ilParSimi1.

INSDC accession	Name	Length (Mb)	GC%
OZ194413.1	1	23.58	38
OZ194414.1	2	23.35	38
OZ194415.1	3	23.29	38
OZ194416.1	4	22.87	38
OZ194417.1	5	22.67	37.5
OZ194418.1	6	22.29	37.5
OZ194419.1	7	22.07	37.5
OZ194420.1	8	21.94	38
OZ194421.1	9	21.64	38
OZ194422.1	10	21.63	37.5
OZ194423.1	11	21.35	37.5
OZ194424.1	12	20.85	37.5
OZ194425.1	13	20.19	38
OZ194426.1	14	19.96	37.5
OZ194427.1	15	19.25	38

INSDC accession	Name	Length (Mb)	GC%
OZ194428.1	16	19.18	38
OZ194429.1	17	18.78	38
OZ194430.1	18	18.27	38
OZ194431.1	19	17.77	38
OZ194432.1	20	17.75	37.5
OZ194433.1	21	17.66	38
OZ194434.1	22	14.59	38
OZ194435.1	23	14.43	38
OZ194436.1	24	14.37	38.5
OZ194437.1	25	13.59	38
OZ194438.1	26	12.18	38
OZ194439.1	27	10.95	38.5
OZ194440.1	28	10.81	38.5
OZ194441.1	29	10.58	39.5
OZ194442.1	30	10.08	39.5
OZ194412.1	Z	26.48	38
OZ194443.1	MT	0.02	17.5

The blobtoolkit pipeline was developed using nf-core tooling (Ewels *et al.*, 2020) and MultiQC (Ewels *et al.*, 2016), relying on the Conda package manager, the Bioconda initiative (Grüning *et al.*, 2018), the Biocontainers infrastructure (da Veiga Leprevost *et al.*, 2017), as well as the Docker (Merkel, 2014) and Singularity (Kurtzer *et al.*, 2017) containerisation solutions.

Table 4 contains a list of relevant software tool versions and sources.

Wellcome Sanger Institute – Legal and Governance

The materials that have contributed to this genome note have been supplied by a Darwin Tree of Life Partner. The submission of materials by a Darwin Tree of Life Partner is

subject to the ‘**Darwin Tree of Life Project Sampling Code of Practice**’, which can be found in full on the Darwin Tree of Life website [here](#). By agreeing with and signing up to the Sampling Code of Practice, the Darwin Tree of Life Partner agrees they will meet the legal and ethical requirements and standards set out within this document in respect of all samples acquired for, and supplied to, the Darwin Tree of Life Project.

Further, the Wellcome Sanger Institute employs a process whereby due diligence is carried out proportionate to the nature of the materials themselves, and the circumstances under which they have been/are to be collected and provided for use. The purpose of this is to address and mitigate any potential legal and/or ethical implications of receipt and use of the materials

Table 4. Software tools: versions and sources.

Software tool	Version	Source
BLAST	2.14.0	ftp://ftp.ncbi.nlm.nih.gov/blast/executables/blast+/
BlobToolKit	4.3.9	https://github.com/blobtoolkit/blobtoolkit
BUSCO	5.5.0	https://gitlab.com/ezlab/busco
bwa-mem2	2.2.1	https://github.com/bwa-mem2/bwa-mem2
DIAMOND	2.1.8	https://github.com/bbuchfink/diamond
fasta_windows	0.2.4	https://github.com/tolkit/fasta_windows
FastK	666652151335353eef2fcd58880bcef5bc2928e1	https://github.com/thegenemyers/FASTK
Gfastats	1.3.6	https://github.com/vgl-hub/gfastats
Goat CLI	0.2.5	https://github.com/genomehubs/goat-cli
Hifiasm	0.19.8-r603	https://github.com/chhylp123/hifiasm
MerquyFK	d00d98157618f4e8d1a9190026b19b471055b22e	https://github.com/thegenemyers/MERQUERY.FK
Minimap2	2.24-r1122	https://github.com/lh3/minimap2
MitoHiFi	3	https://github.com/marcelauliano/MitoHiFi
MultiQC	1.14, 1.17, and 1.18	https://github.com/MultiQC/MultiQC
Nextflow	23.10.0	https://github.com/nextflow-io/nextflow
PretextView	0.2.5	https://github.com/sanger-tol/PretextView
samtools	1.19.2	https://github.com/samtools/samtools
sanger-tol/ascc	-	https://github.com/sanger-tol/ascc
sanger-tol/blobtoolkit	0.5.1	https://github.com/sanger-tol/blobtoolkit
Seqtk	1.3	https://github.com/lh3/seqtk
Singularity	3.9.0	https://github.com/sylabs/singularity
TreeVal	1.2.0	https://github.com/sanger-tol/treeval
YaHS	1.2a.2	https://github.com/c-zhou/yahs

as part of the research project, and to ensure that in doing so we align with best practice wherever possible. The overarching areas of consideration are:

- Ethical review of provenance and sourcing of the material
- Legality of collection, transfer and use (national and international)

Each transfer of samples is further undertaken according to a Research Collaboration Agreement or Material Transfer Agreement entered into by the Darwin Tree of Life Partner, Genome Research Limited (operating as the Wellcome Sanger Institute), and in some circumstances other Darwin Tree of Life collaborators.

Data availability

European Nucleotide Archive: *Parectropis similaria* (brindled white spot). Accession number PRJEB71300; <https://identifiers.org/ena.embl/PRJEB71300>. The genome sequence is released openly for reuse. The *Parectropis similaria* genome sequencing initiative is part of the Darwin Tree of Life (DTOL) project (PRJEB40665) and Project Psyche (PRJEB71705). All raw sequence data and the assembly have been deposited in INSDC databases. The genome will be annotated using available RNA-Seq data and presented through the [Ensembl](#) pipeline at the European Bioinformatics Institute. Raw data and assembly accession identifiers are reported in [Table 1](#) and [Table 2](#).

Author information

Members of the University of Oxford and Wytham Woods Genome Acquisition Lab are listed here: <https://doi.org/10.5281/zenodo.12157525>.

Members of the Natural History Museum Genome Acquisition Lab are listed here: <https://doi.org/10.5281/zenodo.12159242>.

Members of the Darwin Tree of Life Barcoding collective are listed here: <https://doi.org/10.5281/zenodo.12158331>.

Members of the Wellcome Sanger Institute Tree of Life Management, Samples and Laboratory team are listed here: <https://doi.org/10.5281/zenodo.12162482>.

Members of Wellcome Sanger Institute Scientific Operations: Sequencing Operations are listed here: <https://doi.org/10.5281/zenodo.12165051>.

Members of the Wellcome Sanger Institute Tree of Life Core Informatics team are listed here: <https://doi.org/10.5281/zenodo.12160324>.

Members of the Tree of Life Core Informatics collective are listed here: <https://doi.org/10.5281/zenodo.12205391>.

Members of the Darwin Tree of Life Consortium are listed here: <https://doi.org/10.5281/zenodo.4783558>.

References

- Allio R, Schomaker-Bastos A, Romiguier J, *et al.*: **MitoFinder: efficient automated large-scale extraction of mitogenomic data in target enrichment phylogenomics.** *Mol Ecol Resour.* 2020; **20**(4): 892–905. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Altschul SF, Gish W, Miller W, *et al.*: **Basic Local Alignment Search Tool.** *J Mol Biol.* 1990; **215**(3): 403–410. [PubMed Abstract](#) | [Publisher Full Text](#)
- Bateman A, Martin MJ, Orchard S, *et al.*: **UniProt: the universal protein knowledgebase in 2023.** *Nucleic Acids Res.* 2023; **51**(D1): D523–D531. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Bates A, Clayton-Lucey I, Howard C: **Sanger Tree of Life HMW DNA fragmentation: diagenode Megaruptor³ for LI PacBio.** *protocols.io.* 2023. [Publisher Full Text](#)
- Beasley J, Uhl R, Forrest LL, *et al.*: **DNA barcoding SOPs for the Darwin Tree of Life project.** *protocols.io.* 2023; [Accessed 25 June 2024]. [Publisher Full Text](#)
- Boyes D, Mulhair PO, University of Oxford and Wytham Woods Genome Acquisition Lab, *et al.*: **The genome sequence of the Scarce Umber, *Agriopsis aurantiaria* (Hübner, 1799) [version 1; peer review: 2 approved].** *Wellcome Open Res.* 2023; **8**: 463. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Buchfink B, Reuter K, Drost HG: **Sensitive protein alignments at Tree-of-Life scale using DIAMOND.** *Nat Methods.* 2021; **18**(4): 366–368. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Challis R, Kumar S, Sotero-Caio C, *et al.*: **Genomes on a Tree (GoAT): a versatile, scalable search engine for genomic and sequencing project metadata across the eukaryotic Tree of Life [version 1; peer review: 2 approved].** *Wellcome Open Res.* 2023; **8**: 24. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Challis R, Richards E, Rajan J, *et al.*: **BlobToolKit – interactive quality assessment of genome assemblies.** *G3 (Bethesda).* 2020; **10**(4): 1361–1374. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Cheng H, Concepcion GT, Feng X, *et al.*: **Haplotype-resolved *de novo* assembly using phased assembly graphs with hifiasm.** *Nat Methods.* 2021; **18**(2): 170–175. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Crowley L, Allen H, Barnes I, *et al.*: **A sampling strategy for genome sequencing the British terrestrial arthropod fauna [version 1; peer review: 2 approved].** *Wellcome Open Res.* 2023; **8**: 123. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- da Veiga Leprevost F, Grüning BA, Alves Aflitos S, *et al.*: **BioContainers: an open-source and community-driven framework for software standardization.** *Bioinformatics.* 2017; **33**(16): 2580–2582. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Danecek P, Bonfield JK, Liddle J, *et al.*: **Twelve years of SAMtools and BCFtools.** *GigaScience.* 2021; **10**(2): 2580–2582. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Denton A, Oatley G, Cornwell C, *et al.*: **Sanger Tree of Life sample homogenisation: PowerMash.** *protocols.io.* 2023a. [Publisher Full Text](#)
- Denton A, Yatsenko H, Jay J, *et al.*: **Sanger Tree of Life wet laboratory protocol collection V.1.** *protocols.io.* 2023b. [Publisher Full Text](#)

Diesh C, Stevens GJ, Xie P, *et al.*: **JBrowse 2: a modular genome browser with views of syntenic and structural variation.** *Genome Biol.* 2023; **24**(1): 74. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

do Amaral RJV, Denton A, Yatsenko H, *et al.*: **Sanger Tree of Life RNA extraction: automated MagMax™ mirVana.** *protocols.io.* 2023. [Publisher Full Text](#)

Ewels P, Magnusson M, Lundin S, *et al.*: **MultiQC: summarize analysis results for multiple tools and samples in a single report.** *Bioinformatics.* 2016; **32**(19): 3047–3048. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Ewels PA, Peltzer A, Fillinger S, *et al.*: **The nf-core framework for community-curated bioinformatics pipelines.** *Nat Biotechnol.* 2020; **38**(3): 276–278. [PubMed Abstract](#) | [Publisher Full Text](#)

Formenti G, Abueg L, Brajuka A, *et al.*: **Gfastats: conversion, evaluation and manipulation of genome sequences using assembly graphs.** *Bioinformatics.* 2022; **38**(17): 4214–4216. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

GBIF Secretariat: ***Paractropis similaria* (Hufnagel, 1767).** *GBIF Backbone Taxonomy.* 2023; [Accessed 13 February 2025]. [Reference Source](#)

Grüning B, Dale R, Sjödin A, *et al.*: **Bioconda: sustainable and comprehensive software distribution for the life sciences.** *Nat Methods.* 2018; **15**(7): 475–476. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Harry E: **PretextView (Paired REad TEXTure Viewer): a desktop application for viewing pretext contact maps.** 2022. [Reference Source](#)

Henwood B, Sterling P, Lewington R, *et al.*: **Field guide to the caterpillars of great Britain and Ireland.** London: Bloomsbury, 2020. [Reference Source](#)

Howe K, Chow W, Collins J, *et al.*: **Significantly improving the quality of genome assemblies through curation.** *GigaScience.* 2021; **10**(1): g1aa153. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Jay J, Yatsenko H, Narváez-Gómez JP, *et al.*: **Sanger Tree of Life sample preparation: triage and dissection.** *protocols.io.* 2023. [Publisher Full Text](#)

Kerpedjiev P, Abdennur N, Lekschas F, *et al.*: **HiGlass: web-based visual exploration and analysis of genome interaction maps.** *Genome Biol.* 2018; **19**(1): 125. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Kurtzer GM, Sochat V, Bauer MW: **Singularity: scientific containers for mobility of compute.** *PLoS One.* 2017; **12**(5): e0177459. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Lawniczak MKN, Davey RP, Rajan J, *et al.*: **Specimen and sample metadata standards for biodiversity genomics: a proposal from the Darwin Tree of Life project [version 1; peer review: 2 approved with reservations].** *Wellcome Open Res.* 2022; **7**: 187. [Publisher Full Text](#)

Li H: **Minimap2: pairwise alignment for nucleotide sequences.** *Bioinformatics.* 2018; **34**(18): 3094–3100. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Manni M, Berkeley MR, Seppely M, *et al.*: **BUSCO update: novel and streamlined workflows along with broader and deeper phylogenetic coverage for scoring of eukaryotic, prokaryotic, and viral genomes.** *Mol Biol*

Evol. 2021; **38**(10): 4647–4654.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Merkel D: **Docker: lightweight Linux containers for consistent development and deployment.** *Linux J.* 2014; **2014**(239): 2, [Accessed 2 April 2024]. [Reference Source](#)

Oatley G, Denton A, Howard C: **Sanger Tree of Life HMW DNA extraction: automated MagAttract v.2.** *protocols.io.* 2023. [Publisher Full Text](#)

Pereira L, Sivell O, Sivess L, *et al.*: **DTOL: taxon-specific standard operating procedure for the terrestrial and freshwater arthropods working group.** 2022. [Publisher Full Text](#)

Pointon DL, Eagles W, Sims Y, *et al.*: **sanger-tol/treeval v1.0.0 – Ancient Atlantis.** 2023. [Publisher Full Text](#)

Ranallo-Benavidez TR, Jaron KS, Schatz MC: **GenomeScope 2.0 and Smudgeplot for reference-free profiling of polyploid genomes.** *Nat Commun.* 2020; **11**(1): 1432. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Randle Z, Evans-Hill LJ, Parsons MS, *et al.*: **Atlas of Britain and Ireland's larger moths.** Newbury: Pisces Publications, 2019. [Reference Source](#)

Rao SSP, Huntley MH, Durand NC, *et al.*: **A 3D map of the human genome at kilobase resolution reveals principles of chromatin looping.** *Cell.* 2014; **159**(7): 1665–1680. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Rhie A, McCarthy SA, Fedrigo O, *et al.*: **Towards complete and error-free genome assemblies of all vertebrate species.** *Nature.* 2021; **592**(7856): 737–746. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Rhie A, Walenz BP, Koren S, *et al.*: **Merquy: reference-free quality, completeness, and phasing assessment for genome assemblies.** *Genome Biol.* 2020; **21**(1): 245. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Strickland M, Cornwell C, Howard C: **Sanger Tree of Life fragmented DNA clean up: manual SPRI.** *protocols.io.* 2023. [Publisher Full Text](#)

Twyford AD, Beasley J, Barnes I, *et al.*: **A DNA barcoding framework for taxonomic verification in the Darwin Tree of Life project [version 1; peer review: 2 approved].** *Wellcome Open Res.* 2024; **9**: 339. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Uliano-Silva M, Ferreira JGRN, Krashenninnikova K, *et al.*: **MitoHiFi: a python pipeline for mitochondrial genome assembly from PacBio high fidelity reads.** *BMC Bioinformatics.* 2023; **24**(1): 288. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Vasimuddin M, Misra S, Li H, *et al.*: **Efficient architecture-aware acceleration of BWA-MEM for multicore systems.** In: *2019 IEEE International Parallel and Distributed Processing Symposium (IPDPS).* IEEE, 2019; 314–324. [Publisher Full Text](#)

Waring P, Townsend M, Lewington R: **Field guide to the moths of Great Britain and Ireland: third edition.** Bloomsbury Wildlife Guides, 2017. [Reference Source](#)

Zhou C, McCarthy SA, Durbin R: **YaHS: yet another Hi-C scaffolding tool.** *Bioinformatics.* 2023; **39**(1): btac808. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Open Peer Review

Current Peer Review Status:    

Version 1

Reviewer Report 08 May 2025

<https://doi.org/10.21956/wellcomeopenres.26460.r122080>

© 2025 Hof A. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Arjen Van 't Hof 

Biology Centre of the Czech Academy of Sciences, Budějovice, Czech Republic

The genome sequence of the Brindled White-spot moth has been added to the ever-growing list of genome assemblies produced by the Darwin Tree of Life Consortium. It is always enjoyable to read the background of the species before diving into the assembly details. In this case, the background is rather short, but this is not an oversight of the authors. The Brindled White-spot moth has not been studied in great detail, and therefore, there are not that many noteworthy facts to report.

The manuscript follows the structure we are used to from Darwin Tree of Life genome assembly papers, except that there is more emphasis on species confirmation by barcoding which is now a standardized procedure. It is reassuring though that the initial species identification was done by the first author based on entomological knowledge. The assembly statistics are outstanding, in line with what we have come to expect from Darwin Tree of Life.

The language is rich with jargon. I am still not entirely sure what 'localised homologous pairs' means to name an example.

Figure 1 shows tubes coded FD20710430 and FD20710431, but these codes are not explained further in the figure legend or the main text, nor are these codes available in NCBI or EBI databases. It is not clear what these tubes are, what they contain, why there are two of them, and why they are in the photograph.

Is the rationale for creating the dataset(s) clearly described?

Yes

Are the protocols appropriate and is the work technically sound?

Yes

Are sufficient details of methods and materials provided to allow replication by others?

Partly

Are the datasets clearly presented in a useable and accessible format?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Lepidoptera genetics

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 07 May 2025

<https://doi.org/10.21956/wellcomeopenres.26460.r122082>

© 2025 Ioannidis P. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Panagiotis Ioannidis 

Foundation for Research & Technology - Hellas, Crete, Greece

This paper describes the genome sequencing and assembly for *Parectropis similaria*, a lepidopteran insect. The assembly quality looks great and the data are publicly available for anyone who is interested in downloading and using them.

I only have a minor comment; In the "Assembly quality metrics" section the authors say "...identified 98.5% of the expected gene set (single = 98.1%, duplicated = 0.5%)". I'd suggest that they change this to "...identified 98.5% of the BUSCOs...", since a "gene set" is a different thing.

Looking at the deposited sequencing data, I noticed that there also exist RNAseq data. Were these used? For example, for gene annotation? Maybe the authors should mention this in the manuscript, because it is an important resource and it shouldn't go unnoticed!

Is the rationale for creating the dataset(s) clearly described?

Yes

Are the protocols appropriate and is the work technically sound?

Yes

Are sufficient details of methods and materials provided to allow replication by others?

Yes

Are the datasets clearly presented in a useable and accessible format?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: insect genomics, bioinformatics

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 07 May 2025

<https://doi.org/10.21956/wellcomeopenres.26460.r122079>

© 2025 Whibley A. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Annabel Whibley 

The University of Auckland, Auckland, New Zealand

Boyes and colleagues present the high quality reference genome assembly of the Brindled White-spot moth, *Parectropis similaria*. The natural history, specimen collection, sequencing and bioinformatics workflows are comprehensively and accurately described, with excellent reporting of metadata using DToL templates. Public accession links are active. RNAseq data was obtained from the same individual used for HiC library construction but an assembly has not been provided at this time. The presented genome assembly is of high quality, achieving 7.C.Q65, with the failure to recover the W chromosome indicating that the karyotype is likely ZO. This is the first Data Note that I have reviewed where the manual curation protocol links are active (as opposed to "in prep") and its is great to see this information now available.

Is the rationale for creating the dataset(s) clearly described?

Yes

Are the protocols appropriate and is the work technically sound?

Yes

Are sufficient details of methods and materials provided to allow replication by others?

Yes

Are the datasets clearly presented in a useable and accessible format?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Genomics, Bioinformatics

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 30 April 2025

<https://doi.org/10.21956/wellcomeopenres.26460.r122078>

© 2025 Despres L. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Laurence Despres 

Laboratoire d'Ecologie Alpine (Ringgold ID: 56837), Grenoble, Auvergne-Rhône-Alpes, France

The authors report on the chromosome-level genome assembly of an individual female specimen of the Brindled White-spot moth *Parectropis similaria* (Geometridae). The initial identification by morphology of the two collected specimens was verified by an additional DNA barcoding process.

The genome sequence is 574.79 megabases in span. Most of the assembly is scaffolded into 31 chromosomal pseudomolecules, including the Z sex chromosome. The Z chromosome was assigned by synteny to *Agriopis aurantiaria* (GCA_914767915.1). This species appears to exhibit the ZO karyotype. The assembly was performed using appropriate methods : Pac Bio HiFi long reads (42-fold coverage) and Hi-C data to refine scaffolding, manual curation of the final assembly, and contamination check, following the analysis pipeline and high standard of the Darwin Tree of Life Project for results presentation.

The mitogenome was also assembled. The quality of the assembly was further assessed by the proportion of complete BUSCO genes recovered, which is very high (>98% of Lepidoptera database BUSCO genes lepidoptera_odb10 set). Gene annotation was not performed, although RNAseq data was produced and made available. Protein-coding predictions would be a useful addition to this high quality genome.

This high quality genome assembly will provide a reference genome for further genomic studies on this geometrid moth.

Is the rationale for creating the dataset(s) clearly described?

Yes

Are the protocols appropriate and is the work technically sound?

Yes

Are sufficient details of methods and materials provided to allow replication by others?

Yes

Are the datasets clearly presented in a useable and accessible format?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: population genomics

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.
