



DATA NOTE

The genome sequence of the cylindrical bark beetle, *Colydium elongatum* (Fabricius, 1787) (Coleoptera: Zopheridae)

[version 1; peer review: 4 approved]

Liam M. Crowley<sup>1</sup>,  
University of Oxford and Wytham Woods Genome Acquisition Lab,  
Darwin Tree of Life Barcoding Collective,  
Wellcome Sanger Institute Tree of Life Management, Samples and Laboratory  
team,  
Wellcome Sanger Institute Scientific Operations: Sequencing Operations,  
Wellcome Sanger Institute Tree of Life Core Informatics team,  
Tree of Life Core Informatics collective, Darwin Tree of Life Consortium

<sup>1</sup>University of Oxford, Oxford, England, UK

**v1** First published: 05 Nov 2025, 10:623  
<https://doi.org/10.12688/wellcomeopenres.25078.1>  
Latest published: 05 Nov 2025, 10:623  
<https://doi.org/10.12688/wellcomeopenres.25078.1>

Abstract

We present a genome assembly from an individual male *Colydium elongatum* (cylindrical bark beetle; Arthropoda; Insecta; Coleoptera; Zopheridae). The genome sequence has a total length of 450.16 megabases. Most of the assembly (99.15%) is scaffolded into 10 chromosomal pseudomolecules, including the X and Y sex chromosomes. The mitochondrial genome has also been assembled, with a length of 16.82 kilobases. Gene annotation of this assembly on Ensembl identified 13 883 protein-coding genes. This assembly was generated as part of the Darwin Tree of Life project, which produces reference genomes for eukaryotic species found in Britain and Ireland.

Keywords

*Colydium elongatum*; cylindrical bark beetle; genome sequence; chromosomal; Coleoptera



This article is included in the Tree of Life gateway.

Open Peer Review

Approval Status ✓✓✓✓

|             | 1                    | 2                    | 3                    | 4                    |
|-------------|----------------------|----------------------|----------------------|----------------------|
| version 1   | ✓                    | ✓                    | ✓                    | ✓                    |
| 05 Nov 2025 | <a href="#">view</a> | <a href="#">view</a> | <a href="#">view</a> | <a href="#">view</a> |

1. **Zhonghua Wei**, China West Normal University, Nanchong, China
2. **Lindsey C Perkin**, United States Department of Agriculture ARS, College Station, USA
3. **Chenyang Cai**, Chinese Academy of Sciences, Beijing, China
4. **Arun Arumugaperumal** , Rajalakshmi Engineering College, Chennai, India

Any reports and responses or comments on the article can be found at the end of the article.

**Corresponding author:** Darwin Tree of Life Consortium ([mark.blaxter@sanger.ac.uk](mailto:mark.blaxter@sanger.ac.uk))

**Author roles:** Crowley LM: Investigation, Resources;

**Competing interests:** No competing interests were disclosed.

**Grant information:** This work was supported by Wellcome through core funding to the Wellcome Sanger Institute (220540) and the Darwin Tree of Life Discretionary Award [218328, <https://doi.org/10.35802/218328>].  
*The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.*

**Copyright:** © 2025 Crowley LM *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**How to cite this article:** Crowley LM, University of Oxford and Wytham Woods Genome Acquisition Lab, Darwin Tree of Life Barcoding Collective *et al.* **The genome sequence of the cylindrical bark beetle, *Colydium elongatum* (Fabricius, 1787) (Coleoptera: Zopheridae) [version 1; peer review: 4 approved]** Wellcome Open Research 2025, 10:623  
<https://doi.org/10.12688/wellcomeopenres.25078.1>

**First published:** 05 Nov 2025, 10:623 <https://doi.org/10.12688/wellcomeopenres.25078.1>

## Species taxonomy

Eukaryota; Opisthokonta; Metazoa; Eumetazoa; Bilateria; Protostomia; Ecdysozoa; Panarthropoda; Arthropoda; Mandibulata; Pancrustacea; Hexapoda; Insecta; Dicondylia; Pterygota; Neoptera; Endopterygota; Coleoptera; Polyphaga; Cucujiformia; Tenebrionoidea; Zopheridae; Colydiinae; *Colydium*; *Colydium elongatum* (Fabricius, 1787) (NCBI:txid295817)

## Background

*Colydium elongatum* (Fabricius, 1787) is a small, slender, cylindrical zopherid (Colydiinae) with a glossy, almost glabrous dorsum and costate elytra. It is morphologically adapted for movement under bark (Węgrzynowicz, 1999).

It is widespread in Europe and also recorded from North Africa (Węgrzynowicz, 1999). Within the western Palearctic it occurs alongside *C. filiforme*, with a third species (*C. noblecourti*) recently added (Parmain *et al.*, 2024). Adults are saproxylic, living under the bark of dead or dying trees in the galleries of other wood-dwelling insects. *C. elongatum* has been reported to feed by facultative predation on eggs or young larvae of xylophages, or as commensal feeders on fungi and decaying organic matter. Here “saproxylic” follows the usage for species dependent on wood-decay processes in living as well as dead trees (Alexander, 2008).

We present a chromosome-level genome sequence for *Colydium elongatum*, the first genome for the genus *Colydium* as of October 2025 (NCBI datasets, O’Leary *et al.*, 2024). The assembly was produced using the Tree of Life pipeline from a specimen collected in Wytham Woods, Oxfordshire, UK (Figure 1).

## Methods

### Sample acquisition and DNA barcoding

The specimen used for genome sequencing was an adult male *Colydium elongatum* (specimen ID Ox002331, ToLID icColElon2; Figure 1), collected from Wytham Woods, Oxfordshire, UK (latitude 51.77, longitude −1.336) on 2022-05-27. A second

specimen collected on the same occasion was used for Hi-C sequencing (specimen ID Ox002330, ToLID icColElon1). The specimens were collected and identified by Liam Crowley. Sample metadata were collected in line with the Darwin Tree of Life project standards described by Lawniczak *et al.* (2022).

The initial identification was verified by an additional DNA barcoding process according to the framework developed by Twyford *et al.* (2024). A small sample was dissected from one of the specimens and stored in ethanol, while the remaining parts were shipped on dry ice to the Wellcome Sanger Institute (WSI) (see the protocol). The tissue was lysed, the COI marker region was amplified by PCR, and amplicons were sequenced and compared to the BOLD database, confirming the species identification (Crowley *et al.*, 2023). Following whole genome sequence generation, the relevant DNA barcode region was also used alongside the initial barcoding data for sample tracking at the WSI (Twyford *et al.*, 2024). The standard operating procedures for Darwin Tree of Life barcoding are available on protocols.io.

### Nucleic acid extraction

Protocols for high molecular weight (HMW) DNA extraction developed at the Wellcome Sanger Institute (WSI) Tree of Life Core Laboratory are available on protocols.io (Howard *et al.*, 2025). The icColElon2 sample was weighed and triaged to determine the appropriate extraction protocol. Tissue from the whole organism was homogenised by powermashing using a PowerMasher II tissue disruptor. HMW DNA was extracted using the Automated MagAttract v2 protocol. We used centrifuge-mediated fragmentation to produce DNA fragments in the 8–10 kb range, following the Covaris g-TUBE protocol for ultra-low input (ULI). Sheared DNA was purified by automated SPRI (solid-phase reversible immobilisation). The concentration of the sheared and purified DNA was assessed using a Nanodrop spectrophotometer and Qubit Fluorometer using the Qubit dsDNA High Sensitivity Assay kit. Fragment size distribution was evaluated by running the sample on the FemtoPulse system. For this sample, the final post-shearing DNA had a Qubit concentration of 1.86 ng/μL and a yield of 241.80 ng.

### PacBio HiFi library preparation and sequencing

Library preparation and sequencing were performed at the WSI Scientific Operations core. Prior to library preparation, the DNA was fragmented to ~10 kb. Ultra-low-input (ULI) libraries were prepared using the PacBio SMRTbell® Express Template Prep Kit 2.0 and gDNA Sample Amplification Kit. Samples were normalised to 20 ng DNA. Single-strand overhang removal, DNA damage repair, and end-repair/A-tailing were performed according to the manufacturer’s instructions, followed by adapter ligation. A 0.85× pre-PCR clean-up was carried out with Promega ProNex beads.

The DNA was evenly divided into two aliquots for dual PCR (reactions A and B), both following the manufacturer’s protocol. A 0.85× post-PCR clean-up was performed with ProNex beads. DNA concentration was measured using a Qubit Fluorometer v4.0 (Thermo Fisher Scientific) with the Qubit



**Figure 1.** Photograph of the *Colydium elongatum* (icColElon2) specimen used for genome sequencing.

HS Assay Kit, and fragment size was assessed on an Agilent Femto Pulse Automated Pulsed Field CE Instrument (Agilent Technologies) using the gDNA 55 kb BAC analysis kit. PCR reactions A and B were then pooled, ensuring a total mass of  $\geq 500$  ng in 47.4  $\mu$ L.

The pooled sample underwent another round of DNA damage repair, end-repair/A-tailing, and hairpin adapter ligation. A 1 $\times$  clean-up was performed with ProNex beads, followed by DNA quantification using the Qubit and fragment size analysis using the Agilent Femto Pulse. Size selection was performed on the Sage Sciences PippinHT system, with target fragment size determined by Femto Pulse analysis (typically 4–9 kb). Size-selected libraries were cleaned with 1.0 $\times$  ProNex beads and normalised to 2 nM before sequencing.

The sample was sequenced on a Revio instrument (Pacific Biosciences). The prepared library was normalised to 2 nM, and 15  $\mu$ L was used for making complexes. Primers were annealed and polymerases bound to generate circularised complexes, following the manufacturer's instructions. Complexes were purified using 1.2X SMRTbell beads, then diluted to the Revio loading concentration (200–300 pM) and spiked with a Revio sequencing internal control. The sample was sequenced on a Revio 25M SMRT cell. The SMRT Link software (Pacific Biosciences), a web-based workflow manager, was used to configure and monitor the run and to carry out primary and secondary data analysis.

## Hi-C

### Sample preparation and crosslinking

The Hi-C sample was prepared from 20–50 mg of frozen whole organism tissue of the icColElon1 sample using the Arima-HiC v2 kit (Arima Genomics). Following the manufacturer's instructions, tissue was fixed and DNA crosslinked using TC buffer to a final formaldehyde concentration of 2%. The tissue was homogenised using the Diagnocine Power Masher-II. Crosslinked DNA was digested with a restriction enzyme master mix, biotinylated, and ligated. Clean-up was performed with SPRISelect beads before library preparation. DNA concentration was measured with the Qubit Fluorometer (Thermo Fisher Scientific) and Qubit HS Assay Kit. The biotinylation percentage was estimated using the Arima-HiC v2 QC beads.

### Hi-C library preparation and sequencing

Biotinylated DNA constructs were fragmented using a Covaris E220 sonicator and size selected to 400–600 bp using SPRISelect beads. DNA was enriched with Arima-HiC v2 kit Enrichment beads. End repair, A-tailing, and adapter ligation were carried out with the NEBNext Ultra II DNA Library Prep Kit (New England Biolabs), following a modified protocol where library preparation occurs while DNA remains bound to the Enrichment beads. Library amplification was performed using KAPA HiFi HotStart mix and a custom Unique Dual Index (UDI) barcode set (Integrated DNA Technologies). Depending on sample concentration and biotinylation percentage determined at the crosslinking stage, libraries were amplified with 10–16 PCR cycles. Post-PCR clean-up was performed

with SPRISelect beads. Libraries were quantified using the AccuClear Ultra High Sensitivity dsDNA Standards Assay Kit (Biotium) and a FLUOstar Omega plate reader (BMG Labtech).

Prior to sequencing, libraries were normalised to 10 ng/ $\mu$ L. Normalised libraries were quantified again and equimolar and/or weighted 2.8 nM pools were created. Pool concentrations were checked using the Agilent 4200 TapeStation (Agilent) with High Sensitivity D500 reagents before sequencing. Sequencing was performed using paired-end 150 bp reads on the Illumina NovaSeq 6000.

## Genome assembly

Prior to assembly of the PacBio HiFi reads, a database of  $k$ -mer counts ( $k = 31$ ) was generated from the filtered reads using FastK. GenomeScope2 (Ranallo-Benavidez *et al.*, 2020) was used to analyse the  $k$ -mer frequency distributions, providing estimates of genome size, heterozygosity, and repeat content.

The HiFi reads were assembled using Hifiasm (Cheng *et al.*, 2021) with the --primary option. The Hi-C reads (Rao *et al.*, 2014) were mapped to the primary contigs using bwa-mem2 (Vasimuddin *et al.*, 2019), and the contigs were scaffolded in YaHS (Zhou *et al.*, 2023) with the --break option for handling potential misassemblies. The scaffolded assemblies were evaluated using Gfastats (Formenti *et al.*, 2022), BUSCO (Manni *et al.*, 2021) and MERQUERY.FK (Rhie *et al.*, 2020). The organelle genomes were assembled using MitoHiFi (Uliano-Silva *et al.*, 2023).

## Assembly curation

The assembly was decontaminated using the Assembly Screen for Cobionts and Contaminants (ASCC) pipeline. TreeVal was used to generate the flat files and maps for use in curation. Manual curation was conducted primarily in PretextView and HiGlass (Kerpedjiev *et al.*, 2018). Scaffolds were visually inspected and corrected as described by Howe *et al.* (2021). Manual corrections included 17 breaks and 73 joins. This reduced the scaffold count by 19.7% and increased the scaffold N50 by 98.7%. The curation process is described at <https://gitlab.com/wtsi-grit/rapid-curation>. PretextViewSnapshot was used to generate a Hi-C contact map of the final assembly.

## Assembly quality assessment

The Merquery.FK tool (Rhie *et al.*, 2020) was run in a Singularity container (Kurtzer *et al.*, 2017) to evaluate  $k$ -mer completeness and assembly quality for the primary and alternate haplotypes using the  $k$ -mer databases ( $k = 31$ ) computed prior to genome assembly. The analysis outputs included assembly QV scores and completeness statistics.


The genome was analysed using the BlobToolKit pipeline, a Nextflow implementation of the earlier Snakemake version (Challis *et al.*, 2020). The pipeline aligns PacBio reads using minimap2 (Li, 2018) and SAMtools (Danecek *et al.*, 2021) to generate coverage tracks. It runs BUSCO (Manni *et al.*, 2021) using lineages identified from the NCBI Taxonomy (Schoch *et al.*, 2020). For the three domain-level lineages, BUSCO

genes are aligned to the UniProt Reference Proteomes database (Bateman *et al.*, 2023) using DIAMOND blastp (Buchfink *et al.*, 2021). The genome is divided into chunks based on the density of BUSCO genes from the closest taxonomic lineage, and each chunk is aligned to the UniProt Reference Proteomes database with DIAMOND blastx. Sequences without hits are chunked using seqtk and aligned to the NT database with blastn (Altschul *et al.*, 1990). The BlobToolKit suite consolidates all outputs into a blobdir for visualisation. The BlobToolKit pipeline was developed using nf-core tooling (Ewels *et al.*, 2020) and MultiQC (Ewels *et al.*, 2016), with containerisation through Docker (Merkel, 2014) and Singularity (Kurtzer *et al.*, 2017).

Genome sequence report

Sequence data

PacBio sequencing of the *Colydium elongatum* specimen generated 21.96 Gb (gigabases) from 2.64 million reads, which were used to assemble the genome. GenomeScope2.0 analysis estimated the haploid genome size at 420.74 Mb, with a heterozygosity of 0.68% and repeat content of 32.45% (Figure 2). These estimates guided expectations for the assembly. Based on the estimated genome size, the sequencing data provided approximately 50x coverage. Hi-C sequencing produced 87.90 Gb from 582.09 million reads, which were used to scaffold the assembly. Table 1 summarises the specimen and sequencing details.



**Figure 2. Frequency distribution of *k*-mers generated using GenomeScope2.** The plot shows observed and modelled *k*-mer spectra, providing estimates of genome size, heterozygosity, and repeat content based on unassembled sequencing reads.

**Table 1. Specimen and sequencing data for BioProject PRJEB74681.**

| Platform                      | PacBio HiFi              | Hi-C                  |
|-------------------------------|--------------------------|-----------------------|
| ToLID                         | icColElon2               | icColElon1            |
| Specimen ID                   | Ox002331                 | Ox002330              |
| BioSample (source individual) | SAMEA112232558           | SAMEA112232557        |
| BioSample (tissue)            | SAMEA112233006           | SAMEA112233005        |
| Tissue                        | whole organism           | whole organism        |
| Instrument                    | Revio                    | Illumina NovaSeq 6000 |
| Run accessions                | ERR12875211; ERR12875212 | ERR12893017           |
| Read count total              | 2.64 million             | 582.09 million        |
| Base count total              | 21.96 Gb                 | 87.90 Gb              |



Assembly statistics

The primary haplotype was assembled, and contigs corresponding to an alternate haplotype were also deposited in INSDC databases. The final assembly has a total length of 450.16 Mb in 162 scaffolds, with 322 gaps, and a scaffold N50 of 52.95 Mb (Table 2).

Table 2. Genome assembly statistics.

|                               |                         |
|-------------------------------|-------------------------|
| Assembly name                 | icColElon2.1            |
| Assembly accession            | GCA_964656385.1         |
| Alternate haplotype accession | GCA_964656365.1         |
| Assembly level                | chromosome              |
| Span (Mb)                     | 450.16                  |
| Number of chromosomes         | 10                      |
| Number of contigs             | 484                     |
| Contig N50                    | 2.41 Mb                 |
| Number of scaffolds           | 162                     |
| Scaffold N50                  | 52.95 Mb                |
| Sex chromosomes               | X and Y                 |
| Organelles                    | Mitochondrion: 16.82 kb |

Most of the assembly sequence (99.15%) was assigned to 10 chromosomal-level scaffolds, representing 8 autosomes and the X and Y sex chromosomes. These chromosome-level scaffolds, confirmed by Hi-C data, are named according to size (Figure 3; Table 3). Chromosome X was identified by read coverage and alignment to the genome of *Tarphius canariensis* (GCA\_964277315.1). Chromosome Y was identified by read coverage.

The mitochondrial genome was also assembled (length 16.82 kb, OZ210027.1). This sequence is included as a contig in the multifasta file of the genome submission and as a standalone record.

The combined primary and alternate assemblies achieve an estimated QV of 60.8. The *k*-mer completeness is 89.40% for the primary assembly, 68.72% for the alternate haplotype, and 99.52% for the combined assemblies (Figure 4).

BUSCO v.5.5.0 analysis using the endopterygota\_odb10 reference set (*n* = 2 124) identified 99.7% of the expected gene set (single = 98.7%, duplicated = 0.9%). The snail plot in Figure 5 summarises the scaffold length distribution and other assembly statistics for the primary assembly. The blob plot in Figure 6 shows the distribution of scaffolds by GC proportion and coverage.

Table 4 lists the assembly metric benchmarks adapted from Rhie *et al.* (2021) and the Earth BioGenome Project Report on




Figure 3. Hi-C contact map of the *Colydidium elongatum* genome assembly. Assembled chromosomes are shown in order of size and labelled along the axes, with a megabase scale shown below. The plot was generated using PretextSnapshot.

Assembly Standards [September 2024](#). The EBP metric, calculated for the primary assembly, is **6.C.Q63**, meeting the recommended reference standard.

**Table 3. Chromosomal pseudomolecules in the primary genome assembly of *Colydium elongatum* icColElon2.**


| INSDC accession | Molecule | Length (Mb) | GC%   |
|-----------------|----------|-------------|-------|
| OZ210017.1      | 1        | 76.14       | 34.50 |
| OZ210018.1      | 2        | 67.71       | 34.50 |
| OZ210019.1      | 3        | 55.75       | 35    |
| OZ210020.1      | 4        | 52.95       | 35    |
| OZ210021.1      | 5        | 44.93       | 35.50 |
| OZ210022.1      | 6        | 43.04       | 35    |
| OZ210023.1      | 7        | 37.13       | 36    |
| OZ210024.1      | 8        | 37.12       | 35.50 |
| OZ210025.1      | X        | 29          | 35.50 |
| OZ210026.1      | Y        | 2.56        | 34.50 |

Genome annotation report


The *Colydium elongatum* genome assembly (GCA\_964656385.1) was annotated by Ensembl at the European Bioinformatics Institute (EBI). This annotation includes 21 948 transcribed mRNAs from 13 883 protein-coding and 1 841 non-coding genes. The average transcript length is 9 431.64 bp, with an average of 1.40 coding transcripts per gene and 4.88 exons per transcript. For further information about the annotation, please refer to the [Ensembl annotation page](#).

Wellcome Sanger Institute – Legal and Governance

The materials that have contributed to this genome note have been supplied by a Darwin Tree of Life Partner. The submission of materials by a Darwin Tree of Life Partner is subject to the ‘**Darwin Tree of Life Project Sampling Code of Practice**’, which can be found in full on the [Darwin Tree of Life website](#). By agreeing with and signing up to the Sampling Code of Practice, the Darwin Tree of Life Partner agrees they will meet the legal and ethical requirements and standards set out within this document in respect of all samples acquired for, and supplied to, the Darwin Tree of Life Project. Further, the Wellcome Sanger Institute employs a process whereby due diligence is carried out proportionate to the nature of the materials themselves, and the circumstances under which they have been/are to be collected and provided for use. The purpose of this is to address and mitigate any potential legal and/or



**Figure 4. Evaluation of *k*-mer completeness using MerquryFK.** This plot illustrates the recovery of *k*-mers from the original read data in the final assemblies. The horizontal axis represents *k*-mer multiplicity, and the vertical axis shows the number of *k*-mers. The black curve represents *k*-mers that appear in the reads but are not assembled. The green curve corresponds to *k*-mers shared by both haplotypes, and the red and blue curves show *k*-mers found only in one of the haplotypes.



**Figure 5. Assembly metrics for icColElong2.1.** The BlobToolKit snail plot provides an overview of assembly metrics and BUSCO gene completeness. The circumference represents the length of the whole genome sequence, and the main plot is divided into 1 000 bins around the circumference. The outermost blue tracks display the distribution of GC, AT, and N percentages across the bins. Scaffolds are arranged clockwise from longest to shortest and are depicted in dark grey. The longest scaffold is indicated by the red arc, and the deeper orange and pale orange arcs represent the N50 and N90 lengths. A light grey spiral at the centre shows the cumulative scaffold count on a logarithmic scale. A summary of complete, fragmented, duplicated, and missing BUSCO genes in the endopterygota\_odb10 set is presented at the top right. An interactive version of this figure can be accessed on the [BlobToolKit viewer](#).


ethical implications of receipt and use of the materials as part of the research project, and to ensure that in doing so we align with best practice wherever possible. The overarching areas of consideration are:

- Ethical review of provenance and sourcing of the material

- Legality of collection, transfer and use (national and international)

Each transfer of samples is further undertaken according to a Research Collaboration Agreement or Material Transfer Agreement entered into by the Darwin Tree of Life Partner, Genome Research Limited (operating as the Wellcome Sanger





**Figure 6. BlobToolKit GC-coverage plot for *icColElong2.1*.** Blob plot showing sequence coverage (vertical axis) and GC content (horizontal axis). The circles represent scaffolds, with the size proportional to scaffold length and the colour representing phylum membership. The histograms along the axes display the total length of sequences distributed across different levels of coverage and GC content. An interactive version of this figure is available on the [BlobToolKit viewer](#).

**Table 4. Earth Biogenome Project summary metrics for the *Colydium elongatum* assembly.**

| Measure                                        | Value                                                | Benchmark        |
|------------------------------------------------|------------------------------------------------------|------------------|
| EBP summary (primary)                          | 6.C.Q63                                              | 6.C.Q40          |
| Contig N50 length                              | 2.41 Mb                                              | ≥ 1 Mb           |
| Scaffold N50 length                            | 52.95 Mb                                             | = chromosome N50 |
| Consensus quality (QV)                         | Primary: 63.1; alternate: 59.1; combined: 60.8       | ≥ 40             |
| k-mer completeness                             | Primary: 89.40%; alternate: 68.72%; combined: 99.52% | ≥ 95%            |
| BUSCO                                          | C:99.7% [S:98.7%; D:0.9%]; F:0.0%; M:0.3%; n:2 124   | S > 90%; D < 5%  |
| Percentage of assembly assigned to chromosomes | 99.15%                                               | ≥ 90%            |

Institute), and in some circumstances, other Darwin Tree of Life collaborators.

Data availability

European Nucleotide Archive: *Colydium elongatum*. Accession number [PRJEB74681](#). The genome sequence is released openly for reuse. The *Colydium elongatum* genome sequencing initiative is part of the Darwin Tree of Life Project (PRJEB40665) and the Sanger Institute Tree of Life Programme (PRJEB43745). All raw sequence data and the assembly have been deposited in INSDC databases. Raw data and assembly accession identifiers are reported in [Table 1](#) and [Table 2](#).

Production code used in genome assembly at the WSI Tree of Life is available at <https://github.com/sanger-tol>. [Table 5](#) lists software versions used in this study.

Author information

Contributors are listed at the following links:

- Members of the [University of Oxford and Wytham Woods Genome Acquisition Lab](#)
- Members of the [Darwin Tree of Life Barcoding collective](#)
- Members of the [Wellcome Sanger Institute Tree of Life Management, Samples and Laboratory team](#)
- Members of [Wellcome Sanger Institute Scientific Operations – Sequencing Operations](#)
- Members of the [Wellcome Sanger Institute Tree of Life Core Informatics team](#)
- Members of the [Tree of Life Core Informatics collective](#)
- Members of the [Darwin Tree of Life Consortium](#)

Table 5. Software versions and sources.

| Software               | Version             | Source                                                                                                                  |
|------------------------|---------------------|-------------------------------------------------------------------------------------------------------------------------|
| BEDTools               | 2.30.0              | <a href="https://github.com/arq5x/bedtools2">https://github.com/arq5x/bedtools2</a>                                     |
| BLAST                  | 2.14.0              | <a href="ftp://ftp.ncbi.nlm.nih.gov/blast/executables/blast+/">ftp://ftp.ncbi.nlm.nih.gov/blast/executables/blast+/</a> |
| BlobToolKit            | 4.3.9               | <a href="https://github.com/blobtoolkit/blobtoolkit">https://github.com/blobtoolkit/blobtoolkit</a>                     |
| BUSCO                  | 5.5.0               | <a href="https://gitlab.com/ezlab/busco">https://gitlab.com/ezlab/busco</a>                                             |
| bwa-mem2               | 2.2.1               | <a href="https://github.com/bwa-mem2/bwa-mem2">https://github.com/bwa-mem2/bwa-mem2</a>                                 |
| Cooler                 | 0.8.11              | <a href="https://github.com/open2c/cooler">https://github.com/open2c/cooler</a>                                         |
| DIAMOND                | 2.1.8               | <a href="https://github.com/bbuchfink/diamond">https://github.com/bbuchfink/diamond</a>                                 |
| fasta_windows          | 0.2.4               | <a href="https://github.com/tolkit/fasta_windows">https://github.com/tolkit/fasta_windows</a>                           |
| FastK                  | 1.1                 | <a href="https://github.com/thegenemyers/FASTK">https://github.com/thegenemyers/FASTK</a>                               |
| GenomeScope2.0         | 2.0.1               | <a href="https://github.com/tbenavi1/genomescope2.0">https://github.com/tbenavi1/genomescope2.0</a>                     |
| Gfastats               | 1.3.6               | <a href="https://github.com/vgl-hub/gfastats">https://github.com/vgl-hub/gfastats</a>                                   |
| GoaT CLI               | 0.2.5               | <a href="https://github.com/genomehubs/goat-cli">https://github.com/genomehubs/goat-cli</a>                             |
| Hifiasm                | 0.19.8-r603         | <a href="https://github.com/chhylp123/hifiasm">https://github.com/chhylp123/hifiasm</a>                                 |
| HiGlass                | 1.13.4              | <a href="https://github.com/higlass/higlass">https://github.com/higlass/higlass</a>                                     |
| MercuryFK              | 1.1.2               | <a href="https://github.com/thegenemyers/MERQUERY.FK">https://github.com/thegenemyers/MERQUERY.FK</a>                   |
| Minimap2               | 2.24-r1122          | <a href="https://github.com/lh3/minimap2">https://github.com/lh3/minimap2</a>                                           |
| MitoHiFi               | 3                   | <a href="https://github.com/marcelauliano/MitoHiFi">https://github.com/marcelauliano/MitoHiFi</a>                       |
| MultiQC                | 1.14; 1.17 and 1.18 | <a href="https://github.com/MultiQC/MultiQC">https://github.com/MultiQC/MultiQC</a>                                     |
| Nextflow               | 23.10.0             | <a href="https://github.com/nextflow-io/nextflow">https://github.com/nextflow-io/nextflow</a>                           |
| PretextSnapshot        | -                   | <a href="https://github.com/sanger-tol/PretextSnapshot">https://github.com/sanger-tol/PretextSnapshot</a>               |
| PretextView            | 1.0.3               | <a href="https://github.com/sanger-tol/PretextView">https://github.com/sanger-tol/PretextView</a>                       |
| samtools               | 1.19.2              | <a href="https://github.com/samtools/samtools">https://github.com/samtools/samtools</a>                                 |
| sanger-tol/ascc        | 0.1.0               | <a href="https://github.com/sanger-tol/ascc">https://github.com/sanger-tol/ascc</a>                                     |
| sanger-tol/blobtoolkit | 0.6.0               | <a href="https://github.com/sanger-tol/blobtoolkit">https://github.com/sanger-tol/blobtoolkit</a>                       |

| Software                   | Version | Source                                                                                                    |
|----------------------------|---------|-----------------------------------------------------------------------------------------------------------|
| sanger-tol/curationpretext | 1.4.2   | <a href="https://github.com/sanger-tol/curationpretext">https://github.com/sanger-tol/curationpretext</a> |
| Seqtk                      | 1.3     | <a href="https://github.com/lh3/seqtk">https://github.com/lh3/seqtk</a>                                   |
| Singularity                | 3.9.0   | <a href="https://github.com/sylabs/singularity">https://github.com/sylabs/singularity</a>                 |
| TreeVal                    | 1.4.0   | <a href="https://github.com/sanger-tol/treeval">https://github.com/sanger-tol/treeval</a>                 |
| YaHS                       | 1.2a.2  | <a href="https://github.com/c-zhou/yahs">https://github.com/c-zhou/yahs</a>                               |

References

Alexander KNA: **Tree biology and saproxylic Coleoptera: issues of definitions and conservation language.** *Revue d'Ecologie (La Terre et la Vie)*. 2008; (Suppl. 10): 9–13.  
[PubMed Abstract](#) | [Publisher Full Text](#)

Altschul SF, Gish W, Miller W, *et al.*: **Basic Local Alignment Search Tool.** *J Mol Biol*. 1990; **215**(3): 403–410.  
[PubMed Abstract](#) | [Publisher Full Text](#)

Bateman A, Martin MJ, Orchard S, *et al.*: **UniProt: the Universal Protein knowledgebase in 2023.** *Nucleic Acids Res*. 2023; **51**(D1): D523–D531.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Buchfink B, Reuter K, Drost HG: **Sensitive protein alignments at Tree-of-Life scale using DIAMOND.** *Nat Methods*. 2021; **18**(4): 366–368.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Challis R, Richards E, Rajan J, *et al.*: **BlobToolKit – interactive quality assessment of genome assemblies.** *G3 (Bethesda)*. 2020; **10**(4): 1361–1374.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Cheng H, Concepcion GT, Feng X, *et al.*: **Haplotype-resolved *de novo* assembly using phased assembly graphs with hifiasm.** *Nat Methods*. 2021; **18**(2): 170–175.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Crowley L, Allen H, Barnes I, *et al.*: **A sampling strategy for genome sequencing the British terrestrial arthropod fauna [version 1; peer review: 2 approved].** *Wellcome Open Res*. 2023; **8**: 123.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Danecek P, Bonfield JK, Liddle J, *et al.*: **Twelve years of SAMtools and BCFtools.** *GigaScience*. 2021; **10**(2): gfab008.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Ewels P, Magnusson M, Lundin S, *et al.*: **MultiQC: summarize analysis results for multiple tools and samples in a single report.** *Bioinformatics*. 2016; **32**(19): 3047–3048.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Ewels PA, Peltzer A, Fillinger S, *et al.*: **The nf-core framework for community-curated bioinformatics pipelines.** *Nat Biotechnol*. 2020; **38**(3): 276–278.  
[PubMed Abstract](#) | [Publisher Full Text](#)

Formenti G, Abueg L, Brajuka A, *et al.*: **Gfastats: conversion, evaluation and manipulation of genome sequences using assembly graphs.** *Bioinformatics*. 2022; **38**(17): 4214–4216.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Howard C, Denton A, Jackson B, *et al.*: **On the path to reference genomes for all biodiversity: lessons learned and laboratory protocols created in the Sanger Tree of Life core laboratory over the first 2000 species.** *bioRxiv*. 2025.  
[Publisher Full Text](#)

Howe K, Chow W, Collins J, *et al.*: **Significantly improving the quality of genome assemblies through curation.** *GigaScience*. 2021; **10**(1): g1aa153.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Kerpedjiev P, Abdennur N, Lekschas F, *et al.*: **HiGlass: web-based visual exploration and analysis of genome interaction maps.** *Genome Biol*. 2018; **19**(1): 125.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Kurtzer GM, Sochat V, Bauer MW: **Singularity: scientific containers for mobility of compute.** *PLoS One*. 2017; **12**(5): e0177459.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Lawniczak MKN, Davey RP, Rajan J, *et al.*: **Specimen and sample metadata standards for biodiversity genomics: a proposal from the Darwin Tree of Life project [version 1; peer review: 2 approved with reservations].** *Wellcome Open Res*. 2022; **7**: 187.  
[Publisher Full Text](#)

Li H: **Minimap2: pairwise alignment for nucleotide sequences.** *Bioinformatics*. 2018; **34**(18): 3094–3100.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Manni M, Berkeley MR, Seppey M, *et al.*: **BUSCO update: novel and streamlined workflows along with broader and deeper phylogenetic coverage for scoring of eukaryotic, prokaryotic, and viral genomes.** *Mol Biol Evol*. 2021; **38**(10): 4647–4654.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Merkel D: **Docker: lightweight Linux containers for consistent development and deployment.** *Linux J*. 2014; **2014**(239): 2.  
[Reference Source](#)

O’Leary NA, Cox E, Holmes JB, *et al.*: **Exploring and retrieving sequence and metadata for species across the Tree of Life with NCBI datasets.** *Sci Data*. 2024; **11**(1): 732.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Parmain G, Eckelt A, Schuh R: **The genus *Colydium* Fabricius in Europe (Coleoptera, Zopheridae, Colydiinae) with description of a new species, *Colydium noblecourtii* sp. nov.** *Deutsche Entomologische Zeitschrift*. 2024; **71**(2): 289–301.  
[Publisher Full Text](#)

Ranallo-Benavidez TR, Jaron KS, Schatz MC: **GenomeScope 2.0 and Smudgeplot for reference-free profiling of polyploid genomes.** *Nat Commun*. 2020; **11**(1): 1432.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Rao SSP, Huntley MH, Durand NC, *et al.*: **A 3D map of the human genome at kilobase resolution reveals principles of chromatin looping.** *Cell*. 2014; **159**(7): 1665–1680.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Rhie A, McCarthy SA, Fedrigo O, *et al.*: **Towards complete and error-free genome assemblies of all vertebrate species.** *Nature*. 2021; **592**(7856): 737–746.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Rhie A, Walenz BP, Koren S, *et al.*: **Merquy: reference-free quality, completeness, and phasing assessment for genome assemblies.** *Genome Biol*. 2020; **21**(1): 245.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Schoch CL, Ciufo S, Domrachev M, *et al.*: **NCBI taxonomy: a comprehensive update on curation, resources and tools.** *Database (Oxford)*. 2020; **2020**: baaa062.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Twyford AD, Beasley J, Barnes I, *et al.*: **A DNA barcoding framework for taxonomic verification in the Darwin Tree of Life project [version 1; peer review: 2 approved].** *Wellcome Open Res*. 2024; **9**: 339.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Uliano-Silva M, Ferreira JGRN, Krashenninnikova K, *et al.*: **MitoHiFi: a python pipeline for mitochondrial genome assembly from PacBio high fidelity reads.** *BMC Bioinformatics*. 2023; **24**(1): 288.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Vasimuddin M, Misra S, Li H, *et al.*: **Efficient architecture-aware acceleration of BWA-MEM for multicore systems.** In: *2019 IEEE International Parallel and Distributed Processing Symposium (IPDPS)*. IEEE, 2019; 314–324.  
[Publisher Full Text](#)

Węgrzynowicz P: **A revision of the genus *Colydium* Fabricius, 1792 (Coleoptera: Zopheridae: Colydiinae).** *Annales Zoologici*. 1999; **49**(3): 265–328.  
[Reference Source](#)

Zhou C, McCarthy SA, Durbin R: **YaHS: Yet another Hi-C Scaffolding tool.** *Bioinformatics*. 2023; **39**(1): btac808.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

# Open Peer Review

Current Peer Review Status:    

Version 1

Reviewer Report 20 January 2026

<https://doi.org/10.21956/wellcomeopenres.27645.r141907>

© 2026 Arumugaperumal A. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



**Arun Arumugaperumal** 

Department of Biotechnology, Rajalakshmi Engineering College, Chennai, Tamil Nadu, India

The genome note describes the genome sequencing of the cylindrical bark beetle *Colydium elongatum*. The authors have used a male specimen for the sequencing project. They have used PacBio long-read sequencing and Hi-C sequencing data to arrive at an assembly of size 450.16 Mb. Since a male had been used for sequencing, they have obtained the X and Y chromosomes. The mitogenome has also been assembled. They have identified a total of 13,883 protein-coding genes in the assembly. The quality of the assembly is evident from the contig N50 value of 2.41 Mb. BUSCO analysis has also shown a completeness of 99.7% with respect to the endopterygota\_odb10 dataset.

The QV value has been estimated as 60.8, but the EBP metric mentioned is 6.C.Q63. The grant information has an url which appears as text itself.

**Is the rationale for creating the dataset(s) clearly described?**

Yes

**Are the protocols appropriate and is the work technically sound?**

Yes

**Are sufficient details of methods and materials provided to allow replication by others?**

Yes

**Are the datasets clearly presented in a useable and accessible format?**

Yes

**Competing Interests:** No competing interests were disclosed.

**Reviewer Expertise:** Bioinformatics; Genomics

**I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.**

Reviewer Report 08 January 2026

<https://doi.org/10.21956/wellcomeopenres.27645.r141904>

© 2026 Cai C. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



**Chenyang Cai**

Chinese Academy of Sciences, Beijing, Beijing, China

This manuscript presents a high-quality, chromosome-level genome assembly for *Colydium elongatum*, representing the first genomic resource for the genus *Colydium* and one of the few assemblies available for the zopherid subfamily Colydiinae. The study follows the standard Darwin Tree of Life pipeline and provides extensive methodological detail, strong assembly metrics, and a well-documented workflow from sampling to quality assessment. The dataset will clearly be a valuable reference for coleopteran comparative genomics, systematics, and studies of saproxylic insects. the manuscript is clearly written, well structured, and methodologically sound. As this represents the first genomic resource for the genus, it would strengthen the manuscript to expand briefly on: 1) The phylogenetic position of Zopheridae within Tenebrionoidea and 2) What evolutionary questions this genome may help address (e.g., adaptations to saproxylic habitats, body elongation, feeding strategies).

**Is the rationale for creating the dataset(s) clearly described?**

Yes

**Are the protocols appropriate and is the work technically sound?**

Yes

**Are sufficient details of methods and materials provided to allow replication by others?**

Yes

**Are the datasets clearly presented in a useable and accessible format?**

Yes

**Competing Interests:** No competing interests were disclosed.

**Reviewer Expertise:** beetles; phylogenomics; genomes

**I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.**



Reviewer Report 05 January 2026

<https://doi.org/10.21956/wellcomeopenres.27645.r141909>

© 2026 Perkin L. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



**Lindsey C Perkin**

SPARC, ICCDRU, United States Department of Agriculture ARS, College Station, TX, USA

This report is well written. I only have a couple comments.

rationale: while I agree everything should be sequenced without question, it would be nice to learn why it is important to sequence *C. elongatum*. Pest? Important to the ecosystem? This was partially described but more would be nice.

methods: was the second specimen used for Hi-C also male?

**Is the rationale for creating the dataset(s) clearly described?**

Partly

**Are the protocols appropriate and is the work technically sound?**

Yes

**Are sufficient details of methods and materials provided to allow replication by others?**

Yes

**Are the datasets clearly presented in a useable and accessible format?**

Yes

**Competing Interests:** No competing interests were disclosed.

**Reviewer Expertise:** weevil genomics

**I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.**

Reviewer Report 21 November 2025

<https://doi.org/10.21956/wellcomeopenres.27645.r138748>

© 2025 Wei Z. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**Zhonghua Wei**

China West Normal University, Nanchong, China

This study presents the genome assembly and annotation of *Colydium elongatum*, which holds scientific significance for indexing.

1. Figure 1 was blurry, please provide a clear image for this species.

2.

The genome annotation file should be publicly accessible to allow for the verification of potential issues in the annotation.

3. In the References section, some journals are not abbreviated, such as: *Deutsche Entomologische Zeitschrift* and *Journal of Animal Ecology*. Please do the needful.

**Is the rationale for creating the dataset(s) clearly described?**

Yes

**Are the protocols appropriate and is the work technically sound?**

Yes

**Are sufficient details of methods and materials provided to allow replication by others?**

Yes

**Are the datasets clearly presented in a useable and accessible format?**

Yes

**Competing Interests:** No competing interests were disclosed.

**I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.**